

SURVIVIN-INTERACTING PROTEINS AND USE THEREOF

Related U.S. Applications

This application claims priority to U.S. Provisional Application Serial No.
5 60/276,179 filed on March 15, 2001, U.S. Provisional Application Serial No. 60/307,233
filed on July 23, 2001, and U.S. Provisional Application filed on October 25, 2001, all of
which are incorporated herein by reference in their entirety.

Field of the Invention

10 The present invention generally relates to protein-protein interactions, particularly
to protein complexes formed by protein-protein interactions and methods of use thereof.

Background of the Invention

The prolific output from numerous genomic sequencing efforts, including the
15 Human Genome Project, is creating an ever-expanding foundation for large-scale study
of protein function. Indeed, this emerging field of proteomics can appropriately be
viewed as a bridge that connects DNA sequence information to the physiology and
pathology of intact organisms. As such, proteomics – the large-scale study of protein
function – will likely be starting point for the development of many future
20 pharmaceuticals. The efficiency of drug development will therefore depend on the
diversity and robustness of the methods used to elucidate protein function, i.e., the
proteomic tools that are available.

Several approaches are generally known in the art for studying protein function.
One method is to analyze the DNA sequence of a particular gene and the amino acid
25 sequence coded by the gene in the context of sequences of genes with known functions.
Generally, similar functions can be predicted based on sequence homologies. This
“homology method” has been widely used, and powerful computer programs have been

designed to facilitate homology analysis. *See, e.g., Altschul et al., Nucleic Acids Res.*, 25:3389-3402 (1997). However, this method is useful only when the function of a homologous protein is known.

Another useful approach is to interfere with the expression of a particular gene in a cell or organism and examine the consequent phenotypic effects. For example, Fire *et al., Nature*, 391:806-811 (1998) disclose an "RNA interference" assay in which double-stranded RNA transcripts corresponding to a particular target gene are injected into cells or organisms to determine the phenotype associated with the disrupted expression of that gene. Alternatively, transgenic technologies can be utilized to delete or "knock out" a particular gene in an organism and the effect of the gene knockout is determined. *See e.g., Winzeler et al., Science*, 285:901-906 (1999); Zambrowicz *et al., Nature*, 392:608-611 (1998). The phenotypic effects resulting from the disruption of expression of a particular gene can shed some light on the functions of the gene. However, the techniques involved are complex and the time required for a phenotype to appear can be long, especially in mammals. In addition, in many cases disruption of a particular gene may not cause any detectable phenotypic effect.

Gene functions can also be uncovered by genetic linkage analysis. For example, genes responsible for certain diseases may be identified by positional cloning. Alternatively, gene function may be inferred by comparing genetic variations among individuals in a population and correlating particular phenotypes with the genetic variations. Such linkage analyses are powerful tools, particularly when genetic variations exist in a traceable population from which samples are readily obtainable. However, readily identifiable genetic diseases are rare and samples from a large population with genetic variations are not easily accessible. In addition, it is also possible that a gene identified in a linkage analysis does not contribute to the associated disease or symptom but rather is simply linked to unknown genetic variations that cause the phenotypic defects.

With the advance of bioinformatics and publication of the full genome sequence of many organisms, computational methods have also been developed to assign protein functions by comparative genome analysis. For example, Pellegrini *et al., Proc. Natl. Acad. Sci. USA* 96:4285-4288 (1999) discloses a method that constructs a "phylogenetic

profile” that summarizes the presence or absence of a particular protein across a number of organisms as determined by analyzing the genome sequences of the organisms. A protein’s function is predicted to be linked to another protein’s function if the two proteins share the same phylogenetic profile. Another method, the Rosetta Stone method, is based on the theory that separate proteins in one organism are often expressed as separate domains of a fusion protein in another organism. Because the separate domains in the fusion protein are predictably associated with the same function, it can be reasonably predicted that the separate proteins are associated with same functions. Therefore, by discovering separate proteins corresponding to a fusion protein, i.e., the “Rosetta Stone sequence,” functional linkage between proteins can be established. See Marcotte *et al.*, *Science*, 285:751-753 (1999); Enright *et al.*, *Nature*, 402:86-90 (1999). Another computational method is the “gene neighbor method.” See Dandekar *et al.*, *Trends Biochem. Sci.*, 23:324-328 (1998); Overbeek *et al.*, *Proc. Natl. Acad. Sci. USA* 96:2896-2901 (1999). This method is based on the likelihood that if two genes are found to be neighbors in several different genomes, the proteins encoded by the genes share a common function.

While the methods described above are useful in analyzing protein functions, they are constrained by various practical limitations such as unavailability of suitable samples, inefficient assay procedures, and limited reliability. The computational methods are useful in linking proteins by function. However, they are only applicable to certain proteins, and the linkage maps established therewith are sketchy. That is, the maps lack specific information that describes how proteins function in relation to each other within the functional network. Indeed, none of the methods places the identified protein functions in the context of protein-protein interactions.

In contrast with the traditional view of protein function, which focuses on the action of a single protein molecule, a modern expanded view of protein function defines a protein as an element in an interaction network. See Eisenberg *et al.*, *Nature*, 405:823-826 (2000). That is, a full understanding of the functions of a protein will require knowledge of not only the characteristics of the protein itself, but also its interactions or connections with other proteins in the same interacting network. In essence, protein-protein interactions form the basis of almost all biological processes, and each biological

process is composed of a network of interacting proteins. For example, cellular structures such as cytoskeletons, nuclear pores, centrosomes, and kinetochores are formed by complex interactions among a multitude of proteins. Many enzymatic reactions are associated with large protein complexes formed by interactions among enzymes, protein substrates, and protein modulators. In addition, protein-protein interactions are also part of the mechanisms for signal transduction and other basic cellular functions such as DNA replication, transcription, and translation. For example, the complex transcription initiation process generally requires protein-protein interactions among numerous transcription factors, RNA polymerase, and other proteins. *See e.g.*, Tjian and Maniatis, *Cell*, 77:5-8 (1994).

Because most proteins function through their interactions with other proteins, if a test protein interacts with a known protein, one can reasonably predict that the test protein is associated with the functions of the known protein, e.g., in the same cellular structure or same cellular process as the known protein. Thus, interaction partners can provide an immediate and reliable understanding towards the functions of the interacting proteins. By identifying interacting proteins, a better understanding of disease pathways and the cellular processes that result in diseases may be achieved, and important regulators and potential drug targets in disease pathways can be identified.

There has been much interest in protein-protein interactions in the field of proteomics. A number of biochemical approaches have been used to identify interacting proteins. These approaches generally employ the affinities between interacting proteins to isolate proteins in a bound state. Examples of such methods include coimmunoprecipitation and copurification, optionally combined with cross-linking to stabilize the binding. Identities of the isolated protein interacting partners can be characterized by, e.g., mass spectrometry. *See e.g.*, Rout *et al.*, *J. Cell. Biol.*, 148:635-651 (2000); Houry *et al.*, *Nature*, 402:147-154 (1999); Winter *et al.*, *Curr. Biol.*, 7:517-529 (1997). A popular approach useful in large-scale screening is the phage display method, in which filamentous bacteriophage particles are made by recombinant DNA technologies to express a peptide or protein of interest fused to a capsid or coat protein of the bacteriophage. A whole library of peptides or proteins of interest can be expressed and a bait protein can be used to screening the library to identify peptides or proteins

capable of binding to the bait protein. *See e.g.*, U.S. Patent Nos. 5,223,409; 5,403,484; 5,571,698; and 5,837,500. Notably, the phage display method only identifies those proteins capable of interacting in an *in vitro* environment, while the coimmunoprecipitation and copurification methods are not amenable to high throughput screening.

The yeast two-hybrid system is a genetic method that overcomes certain shortcomings of the above approaches. The yeast two-hybrid system has proven to be a powerful method for the discovery of specific protein interactions *in vivo*. *See generally*, Bartel and Fields, eds., *The Yeast Two-Hybrid System*, Oxford University Press, New York, NY, 1997. The yeast two-hybrid technique is based on the fact that the DNA-binding domain and the transcriptional activation domain of a transcriptional activator contained in different fusion proteins can still activate gene transcription when they are brought into proximity to each other. In a yeast two-hybrid system, two fusion proteins are expressed in yeast cells. One has a DNA-binding domain of a transcriptional activator fused to a test protein. The other, on the other hand, includes a transcriptional activating domain of the transcriptional activator fused to another test protein. If the two test proteins interact with each other *in vivo*, the two domains of the transcriptional activator are brought together reconstituting the transcriptional activator and activating a reporter gene controlled by the transcriptional activator. *See, e.g.*, U.S. Patent No. 5,283,173.

Because of its simplicity, efficiency and reliability, the yeast two-hybrid system has gained tremendous popularity in many areas of research. In addition, yeast cells are eukaryotic cells. The interactions between mammalian proteins detected in the yeast two-hybrid system typically are bona fide interactions that occur in mammalian cells under physiological conditions. As a matter of fact, numerous mammalian protein-protein interactions have been identified using the yeast two-hybrid system. The identified proteins have contributed significantly to the understanding of many signal transduction pathways and other biological processes. For example, the yeast two-hybrid system has been successfully employed in identifying a large number of novel mammalian cell cycle regulators that are important in complex cell cycle regulations. Using known proteins that are important in cell cycle regulation as baits, other proteins

involved in cell cycle control were identified by virtue of their ability to interact with the baits. *See generally*, Hannon *et al.*, in *The Yeast Two-Hybrid System*, Bartel and Fields, eds., pages 183-196, Oxford University Press, New York, NY, 1997. Examples of mammalian cell cycle regulators identified by the yeast two-hybrid system include

5 CDK4/CDK6 inhibitors (e.g., p16, p15, p18 and p19), Rb family members (e.g., p130), Rb phosphatase (e.g., PP1- α 2), Rb-binding transcription factors (e.g., E2F-4 and E2F-5), General CDK inhibitors (e.g., p21 and p27), CAK cyclin (e.g., cyclin H), and CDK Thr161 phosphatase (e.g., KAP and CDII). *See id* at page 192. "[T]he two-hybrid approach promises to be a useful tool in our ongoing quest for new pieces of the cell

10 cycle puzzle." *See id* at page 193.

The yeast two-hybrid system can be employed to identify proteins that interact with a specific known protein involved in a disease pathway, and thus provide valuable understandings of the disease mechanism. The identified proteins and the protein-protein interactions in which they participate are potential targets for use in identifying new

15 drugs for treating the disease.

Summary of the Invention

It has been discovered that survivin specifically interacts with cytoplasmic dynein light chain 1 ("HDLC1"), beta-actin, ATP-dependent DNA helicase II 70 kD subunit

20 ("DNA helicase II"), Beta-prime subunit of coatomer complex ("COPP"), Osteopontin, alt. transcript 1 ("OSTP"), Na⁺/Ca²⁺-exchange protein 1 ("SLC8A1"), Catenin, alpha 2 ("A2-CAT"). The specific interactions between these proteins and survivin suggest that survivin and the survivin-interacting proteins are involved in common biological processes. In addition, the interactions between such survivin-interacting proteins and

25 survivin lead to the formation of protein complexes both *in vitro* and *in vivo* that contain survivin and one or more of the survivin-interacting proteins. The protein complexes formed under physiological conditions can mediate the functions and biological activities of survivin and HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT. For example, they may be involved in apoptosis. Thus, the survivin-interacting proteins

30 and the protein complexes are potential drug targets for the development of drugs useful

in treating or preventing diseases and disorders associated with the survivin-containing protein complexes or a protein member thereof, or with apoptosis.

In accordance with a first aspect of the present invention, isolated protein complexes are provided comprising survivin and one or more survivin-interacting proteins selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT. In addition, homologues, derivatives, and fragments of survivin and of the survivin-interacting proteins may also be used in forming protein complexes. In a specific embodiment, fragments of survivin and the survivin-interacting proteins containing the protein domains responsible for the interaction between survivin and the survivin-interacting proteins are used in forming a protein complex of the present invention. In another embodiment, an interacting protein member in the protein complexes of the present invention is a fusion protein containing survivin or a homologue, derivative, or fragment thereof. A fusion protein containing a protein selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT or a homologue, derivative, or fragment thereof may also be used in the protein complexes. In yet another embodiment, a protein complex is provided from a hybrid protein, which comprises survivin or a homologue, derivative, or fragment thereof covalently linked, directly or through a linker, to a survivin-interacting protein selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT or a homologue, derivative, or fragment thereof. In addition, nucleic acids encoding the hybrid protein are also provided.

In yet another aspect, the present invention also provides a method for making the protein complexes. The method includes the steps of providing the first protein and the second protein in the protein complexes of the present invention and contacting said first protein with said second protein. In addition, the protein complexes can be prepared by isolation or purification from tissues and cells or produced by recombinant expression of their protein members. The protein complexes can be incorporated into a protein microchip or microarray, which are useful in large-scale high throughput screening assays involving the protein complexes.

In accordance with a second aspect of the invention, antibodies are provided that are immunoreactive with a protein complex of the present invention. In one embodiment,

an antibody is selectively immunoreactive with a protein complex of the present invention. In another embodiment, a bifunctional antibody is provided that has two different antigen binding sites, each being specific to a different interacting protein member in a protein complex of the present invention. The antibodies of the present invention can take various forms including polyclonal antibodies, monoclonal antibodies, 5 chimeric antibodies, antibody fragments such as Fv fragments, single-chain Fv fragments (scFv), Fab' fragments, and F(ab')₂ fragments. Preferably, the antibodies are partially or fully humanized antibodies. The antibodies of the present invention can be readily prepared using procedures generally known in the art. For example, recombinant 10 libraries such as phage display libraries and ribosome display libraries may be used to screen for antibodies with desirable specificities. In addition, various mutagenesis techniques such as site-directed mutagenesis and PCR diversification may be used in combination with the screening assays.

The present invention also provides detection methods for determining whether 15 there is any aberration in a patient with respect to a protein complex having survivin and one or more survivin-interacting proteins selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT. In one embodiment, the method comprises detecting an aberrant concentration of the protein complexes of the present invention. Alternatively, the concentrations of one or more interacting protein 20 members (at the protein or cDNA or mRNA level) of a protein complex of the present invention are measured. In addition, the cellular localization, or tissue or organ distribution of a protein complex of the present invention is determined to detect any aberrant localization or distribution of the protein complex. In another embodiment, mutations in one or more interacting protein members of a protein complex of the present 25 invention can be detected. In particular, it is desirable to determine whether the interacting protein members have any mutations that will lead to, or are associated with, changes in the functional activity of the proteins or changes in their binding affinity to other interacting protein members in forming a protein complex of the present invention. In yet another embodiment, the binding constant of the interacting protein members of 30 one or more protein complexes is determined. A kit may be used for conducting the detection methods of the present invention. Typically, the kit contains reagents useful in

any of the above-described embodiments of the detection methods, including, e.g., antibodies specific to a protein complex of the present invention or interacting members thereof, and oligonucleotides selectively hybridizable to the cDNAs or mRNAs encoding one or more interacting protein members of a protein complex. The detection methods
5 may be useful in diagnosing a disease or disorder such as cancer, viral infection, AIDS, ischemia, stroke, autoimmune diseases, neurodegenerative diseases, inflammatory disorders, sepsis, and osteoporosis, staging the disease or disorder, or identifying a predisposition to the disease or disorder.

The present invention also provides screening methods for selecting modulators
10 of a protein complex formed between survivin or a homologue, derivative or fragment thereof and a survivin-interacting protein selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT or a homologue, derivative, or fragment thereof. Screening methods are also provided for selecting modulators of a survivin-interacting protein selected from the group consisting of
15 HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT. The compounds identified in the screening methods of the present invention can be used in modulating the functions or activities of survivin, the survivin-interacting proteins, or the protein complexes of the present invention. They may also be effective in modulating the cellular functions involving survivin, survivin-interacting proteins or survivin-containing
20 protein complexes, and in preventing or ameliorating diseases or disorders such as cancer, viral infection, AIDS, ischemia, stroke, autoimmune diseases, neurodegenerative diseases, inflammatory disorders, sepsis, and osteoporosis.

Thus, test compounds may be screened in *in vitro* binding assays to identify compounds capable of binding a protein complex of the present invention or survivin or a
25 survivin-interacting protein identified in accordance with the present invention or homologues, derivatives or fragments thereof. The assays may include the steps of contacting the protein complex with a test compound and detecting the interaction between the interacting partners. In addition, *in vitro* dissociation assays may also be employed to select compounds capable of dissociating or destabilizing the protein
30 complexes identified in accordance with the present invention. For example, the assays may entail (1) contacting the interacting members of the protein complex with each other

in the presence of a test compound; and (2) detecting the interaction between the interacting members. An *in vitro* screening assay may also be used to identify compounds that trigger or initiate the formation of, or stabilize, a protein complex of the present invention.

5 In preferred embodiments, *in vivo* assays such as yeast two-hybrid assays and various derivatives thereof, preferably reverse two-hybrid assays, are utilized in identifying compounds that interfere with or disrupt protein-protein interactions between survivin or a homologue, derivative or fragment thereof and a survivin-interacting protein or a homologue, derivative or fragment thereof. In addition, systems such as yeast two-
10 hybrid assays are also useful in selecting compounds capable of triggering or initiating, enhancing or stabilizing protein-protein interactions between survivin or a homologue, derivative or fragment thereof and a survivin-interacting protein selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT or a homologue, derivative or fragment thereof.

15 In a specific embodiment, the screening method includes: (a) providing in a host cell a first fusion protein having a first protein which is survivin or a homologue or derivative or fragment thereof, and a second fusion protein having a second protein which is survivin-interacting protein as provided in the present invention, or a homologue or derivative or fragment thereof, wherein a DNA binding domain is fused to one of the first
20 and second proteins while a transcription-activating domain is fused to the other of said first and second proteins; (b) providing in the host cell a reporter gene, wherein the transcription of the reporter gene is determined by the interaction between the first protein and the second protein; (c) allowing the first and second fusion proteins to interact with each other within the host cell in the presence of a test compound; and (d)
25 determining the presence or absence of expression of the reporter gene.

The present invention further relates to a method for providing a compound capable of modulating an interaction between the interacting protein members in the protein complexes of the present invention, which comprises the steps of providing atomic coordinates defining a three-dimensional structure of a protein complex, and
30 designing or selecting compounds capable of interfering with the interaction between said first protein and said second protein based on said atomic coordinates.

In addition, the present invention also provides a method for selecting a compound capable of modulating a protein-protein interaction between survivin and a survivin-interacting protein in a protein complex, which comprises the steps of (1) contacting a test compound with a survivin-interacting protein or a homologue or derivative or fragment thereof, and (2) determining whether said test compound is capable of binding said protein. In a preferred embodiment, the method further includes testing a selected test compound capable of binding said protein for its ability to interfere with a protein-protein interaction between survivin and said protein, and optionally further testing the selected test compound capable of binding said protein for its ability to modulate cellular activities associated with survivin and/or the survivin-interacting protein.

The present invention also relates to a virtual screen method for providing a compound capable of modulating an interaction between the interacting members in the protein complexes of the present invention. The method comprises the steps of providing atomic coordinates defining a three-dimensional structure of survivin, or a survivin-interacting protein, and designing or selecting compounds capable of binding survivin or the survivin-interacting protein based on said atomic coordinates. In a preferred embodiment, the method further includes testing a selected test compound capable of binding the protein target for its ability to interfere with a protein-protein interaction between survivin and the survivin-interacting protein, and optionally further testing the selected test compound capable of binding protein target for its ability to modulate cellular activities associated with survivin and/or the survivin-interacting protein.

The present invention further provides a composition having two expression vectors. One vector contains a nucleic acid encoding survivin or a homologue, derivative or fragment thereof. Another vector contains a survivin-interacting protein or a homologue, derivative or fragment thereof. In addition, an expression vector is also provided containing (1) a first nucleic acid encoding survivin or a homologue, derivative or fragment thereof; and (2) a second nucleic acid encoding a survivin-interacting protein or a homologue, derivative or fragment thereof.

Host cells are also provided comprising the expression vector(s). In addition, the present invention also provides a host cell having two expression cassettes. One

expression cassette includes a promoter operably linked to a nucleic acid encoding survivin or a homologue, derivative or fragment thereof. Another expression cassette includes a promoter operably linked to a nucleic acid encoding survivin-interacting protein or a homologue, derivative or fragment thereof.

5 In specific embodiments of the host cells or expression vectors, one of the two nucleic acids is linked to a nucleic acid encoding a DNA binding domain, and the other is linked to a nucleic acid encoding a transcription-activation domain, whereby two fusion proteins can be encoded.

10 In accordance with yet another aspect of the present invention, methods are provided for modulating the functions and activities of a survivin-containing protein complex of the present invention, or interacting protein members thereof. The methods may be used in treating or preventing diseases and disorders such as cancer, viral infection, AIDS, ischemia, stroke, autoimmune diseases, neurodegenerative diseases, inflammatory disorders, sepsis, and osteoporosis. In one embodiment, the methods
15 comprise reducing the protein complex concentration and/or inhibiting the functional activities of the protein complex. Alternatively, the concentration and/or activity of survivin or one of the survivin-interacting proteins may be reduced or inhibited. Thus, the methods may include administering to a patient an antibody specific to a protein complex or survivin or a survivin-interacting protein, an antisense oligo or ribozyme
20 selectively hybridizable to a gene or mRNA encoding survivin or a survivin-interacting protein, or a compound identified in a screening assay of the present invention. In addition, gene therapy methods may also be used in reducing the expression of the gene(s) encoding survivin and/or a survivin-interacting protein.

25 In another embodiment, the methods for modulating the functions and activities of a survivin-containing protein complex of the present invention or interacting protein members thereof comprises increasing the protein complex concentration and/or activating the functional activities of the protein complex. Alternatively, the concentration and/or activity of one of the survivin-interacting proteins or survivin may be increased. Thus, a particular survivin-containing protein complex, survivin or a
30 survivin-interacting protein of the present invention may be administered directly to a patient. Or, exogenous genes encoding one or more protein members of a survivin-

containing protein complex may be introduced into a patient by gene therapy techniques. In addition, a patient needing treatment or prevention may also be administered with compounds identified in a screening assay of the present invention capable of triggering or initiating, enhancing or stabilizing protein-protein interactions between survivin or a
5 homologue, derivative or fragment thereof and a survivin-interacting protein selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT, or a homologue, derivative or fragment thereof.

The foregoing and other advantages and features of the invention, and the manner in which the same are accomplished, will become more readily apparent upon
10 consideration of the following detailed description of the invention taken in conjunction with the accompanying examples, which illustrate preferred and exemplary embodiments.

Detailed Description of the Invention

1. Definitions

The terms "polypeptide," "protein," and "peptide" are used herein interchangeably to refer to amino acid chains in which the amino acid residues are linked by peptide bonds or modified peptide bonds. The amino acid chains can be of any length of greater than two amino acids. Unless otherwise specified, the terms "polypeptide,"
20 "protein," and "peptide" also encompass various modified forms thereof. Such modified forms may be naturally occurring modified forms or chemically modified forms. Examples of modified forms include, but are not limited to, glycosylated forms, phosphorylated forms, myristoylated forms, palmitoylated forms, ribosylated forms, acetylated forms, ubiquitinated forms, etc. Modifications also include intra-molecular
25 crosslinking and covalent attachment to various moieties such as lipids, flavin, biotin, polyethylene glycol or derivatives thereof, etc. In addition, modifications may also include cyclization, branching and cross-linking. Further, amino acids other than the conventional twenty amino acids encoded by genes may also be included in a polypeptide.

30 As used herein, the term "interacting" or "interaction" means that two protein domains, fragments or complete proteins exhibit sufficient physical affinity to each other

so as to bring the two “interacting” protein domains, fragments or proteins physically close to each other. An extreme case of interaction is the formation of a chemical bond that results in continual and stable proximity of the two entities. Interactions that are based solely on physical affinities, although usually more dynamic than chemically bonded interactions, can be equally effective in co-localizing two proteins. Examples of physical affinities and chemical bonds include but are not limited to, forces caused by electrical charge differences, hydrophobicity, hydrogen bonds, van der Waals force, ionic force, covalent linkages, and combinations thereof. The state of proximity between the interaction domains, fragments, proteins or entities may be transient or permanent, reversible or irreversible. In any event, it is in contrast to and distinguishable from contact caused by natural random movement of two entities. Typically, although not necessarily, an “interaction” is exhibited by the binding between the interaction domains, fragments, proteins, or entities. Examples of interactions include specific interactions between antigen and antibody, ligand and receptor, enzyme and substrate, and the like.

An “interaction” between two protein domains, fragments or complete proteins can be determined by a number of methods. For example, an interaction can be determined by functional assays such as the two-hybrid systems. Protein-protein interactions can also be determined by various biophysical and biochemical approaches based on the affinity binding between the two interacting partners. Such biochemical methods generally known in the art include, but are not limited to, protein affinity chromatography, affinity blotting, immunoprecipitation, and the like. The binding constant for two interacting proteins, which reflects the strength or quality of the interaction, can also be determined using methods known in the art. *See Phizicky and Fields, Microbiol. Rev.*, 59:94-123 (1995).

As used herein, the term “protein complex” means a composite unit that is a combination of two or more proteins formed by interaction between the proteins. Typically, but not necessarily, a “protein complex” is formed by the binding of two or more proteins together through specific non-covalent binding interactions. However, covalent bonds may also be present between the interacting partners. For instance, the two interacting partners can be covalently crosslinked so that the protein complex becomes more stable.

The term “protein fragment” as used herein means a polypeptide that represents a portion of a protein. When a protein fragment exhibits interactions with another protein or protein fragment, the two entities are said to interact through interaction domains that are contained within the entities.

5 As used herein, the term “domain” means a functional portion, segment or region of a protein, or polypeptide. “Interaction domain” refers specifically to a portion, segment or region of a protein, polypeptide or protein fragment that is responsible for the physical affinity of that protein, protein fragment or isolated domain for another protein, protein fragment or isolated domain.

10 The term “isolated” when used in reference to nucleic acids (which include gene sequences) of this invention is intended to mean that a nucleic acid molecule is present in a form other than found in nature in its original environment with respect to its association with other molecules. For example, since a naturally existing chromosome includes a long nucleic acid sequence, an “isolated nucleic acid” as used herein means a
15 nucleic acid molecule having only a portion of the nucleic acid sequence in the chromosome but not one or more other portions present on the same chromosome. Thus, for example, an isolated gene typically includes no more than 5 kb, preferably no more than 2 kb, more preferably no more than 1 kb naturally occurring nucleic acid sequence that immediately flanks the gene in the naturally existing chromosome or genomic DNA.
20 However, it is noted that an “isolated nucleic acid” as used herein is distinct from a clone in a conventional library such as genomic DNA library and cDNA library in that the clones in a library is still in admixture with almost all the other nucleic acids in a chromosome or a cell. An isolated nucleic acid can be in a vector. An isolated nucleic acid can also be part of a composition so long as the composition is substantially different
25 from the nucleic acid’s original natural environment. In this respect, an isolated nucleic acid can be in a semi-purified state, i.e., in a composition having certain natural cellular components, while it is substantially separated from other naturally occurring nucleic acids and can be readily detected and/or assayed by standard molecular biology techniques. Preferably, an “isolated nucleic acid” is separated from at least 50%, more
30 preferably at least 75%, most preferably at least 90% of other naturally occurring nucleic acids.

The term "isolated nucleic acid" embraces "purified nucleic acid" which means a specified nucleic acid is in a substantially homogenous preparation of nucleic acid substantially free of other cellular components, other nucleic acids, viral materials, or culture medium, or chemical precursors or by-products associated with chemical reactions for chemical synthesis of nucleic acids. Typically, a "purified nucleic acid" can be obtained by standard nucleic acid purification methods. In a purified nucleic acid, preferably the specified nucleic acid molecule constitutes at least 75%, preferably at least 85%, and more preferably at least 95% of the total nucleic acids in the preparation. The term "purified nucleic acid" also means nucleic acids prepared from a recombinant host cell (in which the nucleic acids have been recombinantly amplified and/or expressed) or chemically synthesized nucleic acids.

The term "isolated nucleic acid" also encompasses "recombinant nucleic acid" which is used herein to mean a hybrid nucleic acid produced by recombinant DNA technology having the specified nucleic acid molecule covalently linked to one or more nucleic acid molecules that are not the nucleic acids naturally flanking the specified nucleic acid. Typically, such one or more nucleic acid molecules flanking the specified nucleic acid are no more than 50 kb, preferably no more than 25 kb.

The term "isolated polypeptide" as used herein means a polypeptide molecule is present in a form other than found in nature in its original environment with respect to its association with other molecules. Typically, an "isolated polypeptide" is separated from at least 50%, more preferably at least 75%, most preferably at least 90% of other naturally co-existing polypeptides in a cell or organism.

The term "isolated polypeptide" encompasses a "purified polypeptide" which is used herein to mean a specified polypeptide is in a substantially homogenous preparation substantially free of other cellular components, other polypeptides, viral materials, or culture medium, or when the polypeptide is chemically synthesized, chemical precursors or by-products associated with the chemical synthesis. For a purified polypeptide, preferably the specified polypeptide molecule constitutes at least 75%, preferably at least 85%, and more preferably at least 95% of the total polypeptide in the preparation. A "purified polypeptide" can be obtained from natural or recombinant host cells by standard purification techniques, or by chemically synthesis.

The term “isolated polypeptide” also encompasses a “recombinant polypeptide” which is used herein to mean a hybrid polypeptide produced by recombinant DNA technology or chemical synthesis having a specified polypeptide molecule covalently linked to one or more polypeptide molecules that do not naturally flank the specified polypeptide.

As used herein, the term “homologue,” when used in connection with a first native protein or fragment thereof that is discovered, according to the present invention, to interact with a second native protein or fragment thereof, means a polypeptide that exhibits an amino acid sequence homology and/or structural resemblance to the first native interacting protein, or to one of the interacting domains of the first native protein such that it is capable of interacting with the second native protein. Typically, a protein homologue of a native protein may have an amino acid sequence that is at least 50%, preferably at least 75%, more preferably at least 80%, 85%, 86%, 87%, 88% or 89%, even more preferably at least 90%, 91%, 92%, 93% or 94%, and most preferably 95%, 96%, 97%, 98% or 99% identical to the native protein. Examples of homologues may be the ortholog proteins of other species including animals, plants, yeast, bacteria, and the like. Homologues may also be selected by, e.g., mutagenesis in a native protein. For example, homologues may be identified by site-specific mutagenesis in combination with assays for detecting protein-protein interactions, e.g., the yeast two-hybrid system described below, as will be apparent to skilled artisans apprised of the present invention. Other techniques for detecting protein-protein interactions include, e.g., protein affinity chromatography, affinity blotting, in vitro binding assays, and the like.

For the purpose of comparing two different nucleic acid or polypeptide sequences, one sequence (test sequence) may be described to be a specific “percent identical to” another sequence (reference sequence) in the present disclosure. In this respect, when the length of the test sequence is less than 90% of the length of the reference sequence, the percentage identity is determined by the algorithm of Myers and Miller, *Bull. Math. Biol.*, 51:5-37 (1989) and Myers and Miller, *Comput. Appl. Biosci.*, 4(1):11-7 (1988). Specifically, the identity is determined by the ALIGN program, which is available at <http://www2.igh.cnrs.fr> maintained by IGH, Montpellier, FRANCE. The default parameters can be used.

Where the length of the test sequence is at least 90% of the length of the reference sequence, the percentage identity is determined by the algorithm of Karlin and Altschul, *Proc. Natl. Acad. Sci. USA*, 90:5873-77 (1993), which is incorporated into various BLAST programs. Specifically, the percentage identity is determined by the “BLAST 2 Sequences” tool, which is available at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. See Tatusova and Madden, *FEMS Microbiol. Lett.*, 174(2):247-50 (1999). For pairwise DNA-DNA comparison, the BLASTN 2.1.2 program is used with default parameters (Match: 1; Mismatch: -2; Open gap: 5 penalties; extension gap: 2 penalties; gap x_dropoff: 50; expect: 10; and word size: 11, with filter). For pairwise protein-protein sequence comparison, the BLASTP 2.1.2 program is employed using default parameters (Matrix: BLOSUM62; gap open: 11; gap extension: 1; x_dropoff: 15; expect: 10.0; and wordsize: 3, with filter).

The term “derivative,” when used in connection with a first native protein (or fragment thereof) that is discovered, according to the present invention, to interact with a second native protein (or fragment thereof), means a modified form of the first native protein prepared by modifying the side chain groups of the first native protein without changing the amino acid sequence of the first native protein. The modified form, i.e., the derivative should be capable of interacting with the second native protein. Examples of modified forms include glycosylated forms, phosphorylated forms, myristylated forms, ribosylated forms, ubiquitinated forms, and the like. Derivatives also include hybrid or fusion proteins containing a native protein or a fragment thereof. Methods for preparing such derivative forms should be apparent to skilled artisans. The prepared derivatives can be easily tested for their ability to interact with the native interacting partner using techniques known in the art, e.g., protein affinity chromatography, affinity blotting, in vitro binding assays, yeast two-hybrid assays, and the like.

The term “isolated protein complex” means a protein complex present in a composition or environment that is different from that found in nature - in its native or original cellular or body environment. Preferably, an “isolated protein complex” is separated from at least 50%, more preferably at least 75%, most preferably at least 90% of other naturally co-existing cellular or tissue components. Thus, an “isolated protein complex” may also be a naturally existing protein complex in an artificial preparation or a non-native host cell. An “isolated protein complex” may also be a “purified protein

complex”, that is, a substantially purified form in a substantially homogenous preparation substantially free of other cellular components, other polypeptides, viral materials, or culture medium, or, when the protein components in the protein complex are chemically synthesized, free of chemical precursors or by-products associated with the chemical synthesis. A “purified protein complex” typically means a preparation containing preferably at least 75%, more preferably at least 85%, and most preferably at least 95% a particular protein complex. A “purified protein complex” may be obtained from natural or recombinant host cells or other body samples by standard purification techniques, or by chemical synthesis.

The terms “hybrid protein,” “hybrid polypeptide,” “hybrid peptide,” “fusion protein,” “fusion polypeptide,” and “fusion peptide” are used herein interchangeably to mean a non-naturally occurring protein having a specified polypeptide molecule covalently linked to one or more polypeptide molecules that do not naturally link to the specified polypeptide. Thus, a “hybrid protein” may be two naturally occurring proteins or fragments thereof linked together by a covalent linkage. A “hybrid protein” may also be a protein formed by covalently linking two artificial polypeptides together. Typically but not necessarily, the two or more polypeptide molecules are linked or “fused” together by a peptide bond forming a single non-branched polypeptide chain.

The term “antibody” as used herein encompasses both monoclonal and polyclonal antibodies that fall within any antibody classes, e.g., IgG, IgM, IgA, or derivatives thereof. The term “antibody” also includes antibody fragments including, but not limited to, Fab, F(ab')₂, and conjugates of such fragments, and single-chain antibodies comprising an antigen recognition epitope. In addition, the term “antibody” also means humanized antibodies, including partially or fully humanized antibodies. An antibody may be obtained from an animal, or from a hybridoma cell line producing a monoclonal antibody, or obtained from cells or libraries recombinantly expressing a gene encoding a particular antibody.

The term “selectively immunoreactive” as used herein means that an antibody is reactive thus binds to a specific protein or protein complex, but not other similar proteins or fragments or components thereof.

The term “activity” when used in connection with proteins or protein complexes means any physiological or biochemical activities displayed by or associated with a particular protein or protein complex including but not limited to activities exhibited in biological processes and cellular functions, ability to interact with or bind another molecule or a moiety thereof, binding affinity or specificity to certain molecules, in vitro or in vivo stability (e.g., protein degradation rate, or in the case of protein complexes ability to maintain the form of protein complex), antigenicity and immunogenicity, enzymatic activities, etc. Such activities may be detected or assayed by any of a variety of suitable methods as will be apparent to skilled artisans.

The term “compound” as used herein encompasses all types of organic or inorganic molecules, including but not limited proteins, peptides, polysaccharides, lipids, nucleic acids, small organic molecules, inorganic compounds, and derivatives thereof.

As used herein, the term “interaction antagonist” means a compound that interferes with, blocks, disrupts or destabilizes a protein-protein interaction; blocks or interferes with the formation of a protein complex; or destabilizes, disrupts or dissociates an existing protein complex.

The term “interaction agonist” as used herein means a compound that triggers, initiates, propagates, nucleates, or otherwise enhances the formation of a protein-protein interaction; triggers, initiates, propagates, nucleates, or otherwise enhances the formation of a protein complex; or stabilizes an existing protein complex.

Unless otherwise specified, the term “survivin” as used herein means the human survivin protein. The usage for naming other proteins should be similar unless otherwise specified in the present disclosure.

2. Protein Complexes

Novel protein-protein interactions have been discovered and confirmed using yeast two-hybrid systems. In particular, it has been discovered that survivin specifically interacts with proteins including HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT. Specific fragments capable of conferring interacting properties on survivin, and HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT have also been identified, which are summarized in Table 1. The GenBank Reference

Numbers for the cDNA sequences encoding survivin, and the survivin-interacting proteins are noted in Table 1 below.

Table 1: Binding Regions of Survivin and Its Interacting Partners

| Bait Protein | | | Prey Proteins | | | |
|-------------------------------------------------|------------------------|------|----------------------------------------------------------------|------------------------|------------------------|------|
| Name and GenBank Accession No. | Amino Acid Coordinates | | Names | GenBank Accession Nos. | Amino Acid Coordinates | |
| | Start | Stop | | | Start | Stop |
| Survivin (BIRC5) Accession Number U75285 | 89 | 143 | Cytoplasmic dynein light chain 1 (HDLC1) | U32944 | 1 | 90 |
| | 3 | 99 | Cytoplasmic dynein light chain 1 (HDLC1) | U32944 | -20 | 89 |
| | 47 | 143 | Cytoplasmic dynein light chain 1 (HDLC1) | U32944 | -20 | 89 |
| | 3 | 99 | beta-actin (ACTB) | K00790 | 336 | 375 |
| | 3 | 99 | DNA helicase II, ATP-dependent, 70 kD subunit (KU70) | S38729 | 131 | 404 |
| | 47 | 143 | Beta-prime subunit of coatomer complex (COPP) | X70476 | 796 | 906 |
| | 3 | 99 | Osteopontin, alt. transcript 1 (OSTP) | BC007016 | 1 | 56 |
| | 52 | 143 | Na ⁺ /Ca ²⁺ -exchange protein 1 (SLC8A1) | M91368 | 302 | 575 |
| | 52 | 143 | Catenin, alpha 2 (A2-CAT) | M94151 | 1 | 166 |
| | 52 | 143 | Catenin, alpha 2 (A2-CAT) | M94151 | 55 | 487 |

- 5 Note: The negative numbers in the start column of bait and/or prey protein amino acid coordinates refer to amino acid residues encoded within the 5' untranslated region of native cellular mRNAs, i.e., residues encoded upstream of the normal translational start site.

2.1. Cellular Functions and Disease Involvement

Survivin has clearly been demonstrated to function as an anti-apoptotic factor and also appears to play a role in cytokinesis. Olie et al., *Cancer Res.*, 60(11):2805-9 (2000); Chen et al., *Neoplasia*, 2(3):235-41 (2000). Survivin has been shown to inhibit caspase
5 function and to override the mitotic spindle checkpoint. Suzuki et al., *Oncogene*, 19(10):1346-53 (2000); Li et al., *Nature*, 396(6711):580-4 (1998). In addition, it has been shown that survivin has (positive) cell cycle effects that coincide with its movement from the cytoplasm to the nucleus where it interacts with the Cdk4/p16(INK4a) complex. Suzuki et al., *Oncogene*, 19(29):3225-34 (2000). This is followed by phosphorylation of
10 Rb (anti-apoptotic).

Survivin is one of few known proteins that integrate cell cycle progression and programmed cell death, and thus is involved in preserving homeostasis and developmental morphogenesis (reviewed in Altieri et al., *Lab Invest.*, 79:1327-1333 (1999)). Survivin, also known as API4 or BIRC5, is a 142 amino acid protein that
15 contains a single Zn finger of the type found in other IAP (inhibitor of apoptosis) family members that is necessary for apoptosis inhibition, and a C-terminal RING finger thought to mediate protein-protein interaction (Cahill et al., *Nature*, 392:300-303 (1998)). In addition, survivin contains numerous potential phosphorylation sites. Suppression of survivin expression in HeLa cells results in increased apoptosis and inhibition of
20 proliferation (Ambrosini et al., *J. Biol. Chem.*, 273:11177-11182 (1998)). Furthermore, expression of a phosphorylation-defective (T34A) survivin mutant protein in several human melanoma cell lines triggered apoptosis and enhanced sensitivity to the chemotherapeutic drug cisplatin, and either prevented tumor formation or retarded tumor growth in mice (Grossman et al., *Proc. Natl. Acad. Sci. U.S.A.*, 98:635-640 (2001)).

Survivin is expressed during G2/M phase of the cell cycle and associates with microtubules of the mitotic spindle during mitosis. Disruption of the association of survivin with microtubules results in loss of anti-apoptotic activity and an increase in caspase-3 activity during mitosis (Li et al., *Nature*, 396:580-584 (1998). Survivin interacts directly with, and inhibits, both caspase-3 and caspase-7, and therefore it has
30 been proposed that the anti-apoptotic effects of survivin are due to sequestration of these caspases in an inactive form on microtubules by survivin (Shin et al., *Biochemistry*,

40:1117-1123 (2001)). Together, these results suggest that survivin function counteracts a default apoptotic mechanism at the G2/M transition. Overexpression of survivin in a variety of cancers (e.g. adenocarcinoma and high-grade lymphomas; (Cahill *et al.*, *Nature*, 392:300-303 (1998)) may overcome an apoptotic checkpoint and favor aberrant
5 cell cycle progression.

Survivin interacts with dynein light chain 1. A search of the brain library by yeast two-hybrid analysis using as bait a C-terminal fragment (amino acids 89 to 142) of survivin isolated as an interactor cytoplasmic dynein light chain 1 (HDLC1). The bait is the region of survivin C-terminal to the BIR repeat (baculovirus inhibitor of apoptosis
10 repeat), and it contains a coiled coil (aa 99-142), which is found in some structural proteins, such as myosins, and in some DNA-binding proteins as the so-called leucine-zipper. HDLC1 is an 89-amino acid protein, and the prey isolated here encodes the entire ORF as well as 20 "amino acids" of translated 5'-UTR. Dyneins are molecular motors that translocate along microtubules. Null mutations of *Drosophila dlc1* were lethal and
15 caused embryonic degeneration and widespread apoptotic cell death. Recently, the proapoptotic Bcl-2 family member Bim was shown to be sequestered by LC8 DLC in healthy cells, and this interaction was disrupted by certain apoptotic stimuli. This freed Bim to translocate together with LC8 to Bcl-2 and to neutralize its antiapoptotic activity. Furthermore, it has been found that the 10-kD human DLC1 protein physically interacts
20 with and inhibits the activity of neuronal nitric oxide synthase. Jaffrey and Snyder, *Science*, 274:774-7 (1996). In the brain, nitric oxide is responsible for the glutamate-linked enhancement of 3-prime, 5-prime cyclic guanosine monophosphate levels and may be involved in apoptosis, synaptogenesis, and neuronal development. It is thus not surprising that survivin interacts with a protein having a central role in apoptosis.

25 Searches of the brain library with survivin (aa 3-99 and 47-142) also identified cytoplasmic dynein light chain 1 (HDLC1) as an interactor. The prey is the full-length protein including 19 amino acids of translated 5'-UTR. Survivin is 142 amino acids long, and the previous survivin bait that isolated HDLC1 is amino acids 89-142. HDLC1 either interacts with both the N-terminus (aa 3-99) and C-terminus (aa 89-142) of
30 survivin, or we have fortuitously identified a core HDLC1 binding site on survivin, namely aa 89-99. Survivin may be held in the cytoplasm in a complex with dynein light

chain in the absence of an anti-apoptotic signal. This can occur either appropriately, such as during angiogenesis, or it can occur inappropriately, such as in many cancers. Tran *et al.*, *Biochem. Biophys. Res. Commun.*, 264(3):781-8 (1999); O'Conner *et al.*, *Am. J. Pathol.*, 156(2):393-81 (2000). Pathologically, survivin expression correlates with a bad prognosis for cancer survival. Sarela *et al.*, *Gut*, 46(5):645-50 (2000). Survivin is an example of a more "cancer-specific" drug target, which may be useful in developing anticancer drugs that may have fewer cytotoxic side effects than do current chemotherapeutics. Buolamwini, *Curr. Opin. Chem. Biol.*, 3(4):500-9 (1999).

Survivin is a 142 aa protein that functions as an inhibitor of apoptosis (IAP). Survivin is abundantly expressed in transformed cells of lymphoid and myeloid lineage, adenocarcinoma of the lung, pancreas, colon, breast, and prostate. Survivin expression appears to be developmentally regulated; the protein is expressed in fetal tissues but is absent in adult, terminally differentiated tissues [Ambrosini et al. 1997 Nat Med 3:917]. Expression of survivin, a microtubule-associated protein, is specific to the G2/M phase of the cell cycle.

Survivin acts to suppress apoptosis through inhibition of caspases 3 and 7. Unlike most IAP proteins, survivin contains only a single BIR domain (baculovirus inhibitor of apoptosis repeat) and lacks a RING finger. Survivin has the highest sequence homology to the yeast Bir1 protein that functions in cell division control and chromosome segregation [Li F et al. 2000 J Biol Chem 275:6707; Li F et al. 1998 Nature 396:580].

We have identified an interaction between the C-terminus of survivin (aa89-142) and the full-length 89 aa protein cytoplasmic dynein light chain, HDLC1. The light chain is an accessory subunit of the cytoskeletal motor protein dynein that is responsible for microtubule-associated intracellular movement.

The C-terminal portion of survivin involved in this interaction is downstream of the BIR domain and contains a coiled-coil domain (aa99-142) that appears to be responsible for the interaction of survivin with microtubules. A truncated survivin mutant (M1-G99) binds minimally to polymerized microtubules and is not cytoprotective against taxol-induced apoptosis [Li F et al. 1998 Nature 396:580]. In addition, a point mutant (Cys84Ala) that alters a residue conserved throughout BIR domains, binds indistinguishably from wt survivin to microtubules but is not cytoprotective.

A functional BIR domain and localization to microtubules appear to be necessary for the inhibition of apoptosis by survivin. Inhibiting the interaction between HDLC1 and survivin may result in a loss of survivin from the microtubule and a negation of its function in inhibiting apoptosis. It has been demonstrated that gene targeting of survivin with an antisense mRNA derived from EPR-1 in HeLa cells increased apoptosis and inhibited the growth of these transformed cells [Ambrosini et al. 1998 J Biol Chem 273: 11177].

In support of this approach are data derived with the Bcl-2 family member Bim. The pro-apoptotic activity of Bim appears to be regulated by its interaction with the dynein motor complex through the LC8 cytoplasmic dynein light chain [Puthalakath H 1999 Mol Cell 3:287]. Apoptotic stimuli disrupt the interaction of LC8 with the dynein complex and result in the translocation of LC8 and Bim away from microtubules. It has been hypothesized that this release allows Bim to move to Bcl-2 to inhibit its anti-apoptotic effect.

Survivin interacts with β -actin and DNA helicase II. The search of the brain library with survivin (aa 3-99) identified β -actin and ATP-dependent DNA helicase II, 70 kD subunit (Ku70) as interactors. The β -actin prey isolated here (aa 336-375) is C-terminal. β -actin is a non-muscle cytoskeletal actin involved in cell motility. Survivin and β -actin have not been linked in the literature, however, β -actin mRNA is differentially expressed in cells undergoing apoptosis, suggesting a link between β -actin and apoptosis, and therefore perhaps survivin. Naora and Naora, *Biochem. Biophys. Res. Commun.*, 211(2):491-6 (1995). Actins in general have been linked to apoptosis as targets of caspases. Rossiter et al., *Neuropathol. Appl. Neurobiol.*, 26(4):342-6 (2000). Actins have been shown to undergo rearrangement concomitant with apoptosis. Mashima et al., *Oncogene*, 18(15):2423-30 (1999); Suarez-Huerta et al., *J. Cell Physiol.*, 184(2):239-45 (2000). Interestingly, it has been shown that p53 binds to filamentous actin in a calcium-dependent manner. Metcalfe et al., *Oncogene*, 18(14):2351-5 (1999). Thus, it has been speculated that this may play a role in the transient and reversible nuclear to cytoplasmic shuttling which p53 undergoes. Metcalfe et al., *Oncogene*, 18(14):2351-5 (1999). For instance, during DNA synthesis, when non-pathological DNA strand breaks are present, a p53 response should not be triggered.

Another interactor, Ku70 (aa 131-403), is a single-stranded DNA- and ATP-dependent helicase thought to have a role in DNA repair (DNA damage sensor) and chromosomal translocation (double-strand break repair protein). It has been proposed that the presence or absence of Ku70 determines whether double-stranded breaks are repaired by nonhomologous end joining (DNA damage; Ku70 present) or by homologous recombination (meiosis; Ku70 absent). Goedecke *et al.*, (*Nat. Genet.*, 23(2):194-8 (1999). Li *et al.*, *Mol. Cell*, 2(1):1-8 (1998) suggested that Ku70 is a candidate tumor suppressor gene since mice carrying a disruption of the Ku70 gene showed a propensity for malignant transformation (T-cell lymphomas). Autoantibodies against Ku70 are common in cases of systemic lupus erythematosus and autoimmune thyroiditis (Grave's disease), and this is how Ku70 was first identified. Interestingly, Takeda *et al.*, *J. Immunol.*, 163(11):6269-74 (1999) mentioned that caspase-cleaved proteins can elicit the generation of autoantibodies, because cleavage of self antigens may enhance their immunogenicity. In the context of the survivin-Ku70 interaction, it is possible that in the case of lupus, this represents a pathologic disorder of apoptosis in which Ku70 is inappropriately cleaved. Physiologically, it is likely that survivin plays a role in targeting DNA metabolizing enzymes, such as the helicase Ku70, for proteolysis as one facet of the orchestrated process of programmed cell death. It has been disclosed that an ionizing radiation-induced Ku70-containing complex appears to regulate whether cells undergo apoptosis following a DNA insult. Yang *et al.*, *Proc. Natl. Acad. Sci. U S A*, 97(11):5907-12 (2000). In addition, it has been further suggested that Ku70 up-regulation (following ionization) serves to determine either a course of DNA repair or an arresting response, such as cell death. Brown *et al.*, *J. Biol. Chem.*, 275(9):6651-6 (2000).

The interactions between survivin and proteins including HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT suggest that these proteins are involved in common biological processes including, but not limited to, apoptosis, and disease pathways involving such cellular functions.

2.2. Protein Complexes

Accordingly, the present invention provides protein complexes formed between survivin and one or more survivin-interacting proteins selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT. The present invention also provides protein complexes formed from the interaction between homologues, derivatives or fragments of survivin and one or more of the survivin-interacting proteins in accordance with the present invention. In addition, the present invention further encompasses protein complexes having survivin and homologues, derivatives or fragments of one or more of the survivin-interacting proteins in accordance with the present invention. In yet another embodiment, protein complexes are provided having homologues, derivatives or fragments of survivin and homologues, derivatives or fragments of one or more of the survivin-interacting proteins in accordance with the present invention. In other words, one or more of the interacting protein members of a protein complex of the present invention may be a native protein or a homologue, derivative or fragment of a native protein.

As described above, individual protein fragments involved in the specific protein-protein interactions have been discovered and summarized in Table 1. Accordingly, protein fragments containing the amino acid sequence of the identified regions or homologues or derivatives thereof can be used in forming the protein complexes of the present invention. Fragments capable of interacting with survivin can be provided by the combination of molecular engineering of a nucleic acid encoding a survivin-interacting protein and a method for testing protein-protein interaction. For example, the coordinates in Table 1 can be used as starting points and various fragments of the survivin-interacting protein falling within the coordinates can be generated by deletions from either or both ends of the coordinates. The resulting fragments can be tested for their ability to interact with survivin using any methods known in the art for detecting protein-protein interactions (e.g., yeast two-hybrid method). Likewise, survivin fragments capable of interacting with a survivin-interacting protein, and fragments of other survivin-interacting proteins capable of interacting with survivin can also be identified in a similar manner.

Thus, for example, one interacting partner in a protein complex can be a complete native survivin, a survivin homologue capable of interacting with, e.g., HDLC1, a

survivin derivative, a derivative of the survivin homologue, a survivin fragment capable of interacting with HDLC1 (survivin fragment(s) containing the coordinates shown in Table 1), a derivative of the survivin fragment, or a fusion protein containing (1) complete native survivin, (2) a survivin homologue capable of interacting with HDLC1 or (3) a survivin fragment capable of interacting with HDLC1. Besides native HDLC1, useful interacting partners for survivin or a homologue or derivative or fragment thereof also include homologues of HDLC1 capable of interacting with survivin, derivatives of the native or homologue HDLC1 capable of interacting with survivin, fragments of the HDLC1 capable of interacting with survivin (e.g., a fragment containing the identified interacting regions shown in Table 1), derivatives of the HDLC1 fragments, or fusion proteins containing (1) a complete HDLC1, (2) a HDLC1 homologue capable of interacting with survivin or (3) a HDLC1 fragment capable of interacting with survivin.

HDLC1 fragments capable of interacting with survivin can be the combination of molecular engineering of a HDLC1-encoding nucleic acid and a method for testing protein-protein interaction. For example, the coordinates in Table 1 can be used as starting points and various HDLC1 fragments falling within the coordinates can be generated by deletions from either or both ends of the coordinates. The resulting fragments can be tested for their ability to interact with survivin using any methods known in the art for detecting protein-protein interactions (e.g., yeast two-hybrid method). Likewise, survivin fragments and fragments of other survivin-interacting proteins capable of interacting with HDLC1 can also be identified in a similar manner.

Other protein complexes can be formed in a similar manner based on interactions between survivin and its other interacting partners discovered according to the present invention or homologues, derivatives or fragments of such other interacting partners. In addition, protein complexes containing survivin and two or more members of the group of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT or homologues, derivatives, or fragments thereof can also be formed.

In a specific embodiment of the protein complex of the present invention, two or more interacting partners (survivin and one or more proteins selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT, or homologues, derivatives or fragments thereof) are directly fused together, or covalently

linked together through a peptide linker, forming a hybrid protein having a single unbranched polypeptide chain. Thus, the protein complex may be formed by “intramolecular” interactions between two portions of the hybrid protein. Again, one or both of the fused or linked interacting partners in this protein complex may be a native protein or a homologue, derivative or fragment of a native protein.

As will be apparent to skilled artisans, suitable interacting fragments of a protein capable of interacting with an interacting partner of the protein can be obtained using various techniques well known in the art. For example, since the binding regions of both survivin and HDLC1 are provided in Table 1, various survivin fragments within the binding regions of survivin can be made by chemical synthesis or recombinant DNA technologies. The survivin fragments can then be tested for its ability to interact with HDLC1 using any method known in the art for detecting protein-protein interactions. Examples of such methods include protein affinity chromatography, affinity blotting, in vitro binding assays, yeast two-hybrid assays, and the like.

The protein complexes of the present invention can also be in a modified form. For example, an antibody selectively immunoreactive with the protein complex can be bound to the protein complex. In another example, a non-antibody modulator capable of enhancing the interaction between the interacting partners in the protein complex may be included. Alternatively, the protein members in the protein complex may be cross-linked for purposes of stabilization. Various crosslinking methods may be used. For example, a bifunctional reagent in the form of R-S-S-R' may be used in which the R and R' groups can react with certain amino acid side chains in the protein complex forming covalent linkages. See e.g., Traut *et al.*, in Creighton ed., *Protein Function: A Practical Approach*, IRL Press, Oxford, 1989; Baird *et al.*, *J. Biol. Chem.*, 251:6953-6962 (1976). Other useful crosslinking agents include, e.g., Denny-Jaffee reagent, a heterbiofunctional photoactivable moiety cleavable through an azo linkage (See Denny *et al.*, *Proc. Natl. Acad. Sci. USA*, 81:5286-5290 (1984)), and ¹²⁵I-{S-[N-(3-iodo-4-azidosalicyl)cysteaminy]-2-thiopyridine}, a cysteine-specific photocrosslinking reagent (see Chen *et al.*, *Science*, 265:90-92 (1994)).

The above-described protein complexes may further include any additional components, e.g., other proteins, nucleic acids, lipid molecules, monosaccharides or polysaccharides, ions, etc.

2.3. Methods of Preparing Protein Complexes

The protein complex of the present invention can be prepared by a variety of methods. Specifically, a protein complex can be isolated directly from an animal tissue sample, preferably a human tissue sample containing the protein complex. Alternatively, a protein complex can be purified from host cells that recombinantly express the members of the protein complex. As will be apparent to a skilled artisan, a protein complex can be prepared from a tissue sample or recombinant host cells by coimmunoprecipitation using an antibody immunoreactive with an interacting protein partner, or preferably an antibody selectively immunoreactive with the protein complex as will be discussed in detail below.

The antibodies can be monoclonal or polyclonal. Coimmunoprecipitation is a commonly used method in the art for isolating or detecting bound proteins. In this procedure, generally a serum sample or tissue or cell lysate is admixed with a suitable antibody. The protein complex bound to the antibody is precipitated and washed. The bound protein complexes are then eluted.

Alternatively, immunoaffinity chromatography and immunoblotting techniques may also be used in isolating the protein complexes from native tissue samples or recombinant host cells using an antibody immunoreactive with an interacting protein partner, or preferably an antibody selectively immunoreactive with the protein complex. For example, in protein immunoaffinity chromatography, the antibody is covalently or non-covalently coupled to a matrix (e.g., Sepharose), which is then packed into a column. Extract from a tissue sample, or lysate from recombinant cells is passed through the column where it contacts the antibodies attached to the matrix. The column is then washed with a low-salt solution to wash away the unbound or loosely (non-specifically) bound components. The protein complexes that are retained in the column can be then eluted from the column using a high-salt solution, a competitive antigen of the antibody, a chaotropic solvent, or sodium dodecyl sulfate (SDS), or the like. In immunoblotting,

crude proteins samples from a tissue sample extract or recombinant host cell lysate are fractionated by polyacrylamide gel electrophoresis (PAGE) and then transferred to a membrane, e.g., nitrocellulose. Components of the protein complex can then be located on the membrane and identified by a variety of techniques, e.g., probing with specific antibodies.

In another embodiment, individual interacting protein partners may be isolated or purified independently from tissue samples or recombinant host cells using similar methods as described above. The individual interacting protein partners are then combined under conditions conducive to their interaction thereby forming a protein complex of the present invention. It is noted that different protein-protein interactions may require different conditions. As a starting point, for example, a buffer having 20 mM Tris-HCl, pH 7.0 and 500 mM NaCl may be used. Several different parameters may be varied, including temperature, pH, salt concentration, reducing agent, and the like. Some minor degree of experimentation may be required to determine the optimum incubation condition, this being well within the capability of one skilled in the art once apprised of the present disclosure.

In yet another embodiment, the protein complex of the present invention may be prepared from tissue samples or recombinant host cells or other suitable sources by protein affinity chromatography or affinity blotting. That is, one of the interacting protein partners is used to isolate the other interacting protein partner(s) by binding affinity thus forming protein complexes. Thus, an interacting protein partner prepared by purification from tissue samples or by recombinant expression or chemical synthesis may be bound covalently or non-covalently to a matrix, e.g., Sepharose, which is then packed into a chromatography column. The tissue sample extract or cell lysate from the recombinant cells can then be contacted with the bound protein on the matrix. A low-salt solution is used to wash off the unbound or loosely bound components, and a high-salt solution is then employed to elute the bound protein complexes in the column. In affinity blotting, crude protein samples from a tissue sample or recombinant host cell lysate can be fractionated by polyacrylamide gel electrophoresis (PAGE) and then transferred to a membrane, e.g., nitrocellulose. The purified interacting protein member is then bound to

its interacting protein partner(s) on the membrane forming protein complexes, which are then isolated from the membrane.

It will be apparent to skilled artisans that any recombinant expression methods may be used in the present invention for purposes of expressing the protein complexes or individual interacting proteins. Generally, a nucleic acid encoding an interacting protein member can be introduced into a suitable host cell. For purposes of forming a recombinant protein complex within a host cell, nucleic acids encoding two or more interacting protein members should be introduced into the host cell.

Typically, the nucleic acids, preferably in the form of DNA, are incorporated into a vector to form expression vectors capable of directing the production of the interacting protein member(s) once introduced into a host cell. Many types of vectors can be used for the present invention. Methods for the construction of an expression vector for purposes of this invention should be apparent to skilled artisans apprised of the present disclosure. *See generally, Current Protocols in Molecular Biology*, Vol. 2, Ed. Ausubel, *et al.*, Greene Publish. Assoc. & Wiley Interscience, Ch. 13, 1988; Glover, *DNA Cloning*, Vol. II, IRL Press, Wash., D.C., Ch. 3, 1986; Bitter, *et al.*, in *Methods in Enzymology* 153:516-544 (1987); *The Molecular Biology of the Yeast Saccharomyces*, Eds. Strathern *et al.*, Cold Spring Harbor Press, Vols. I and II, 1982; and Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press, 1989.

Generally, the expression vectors include an expression cassette having a promoter operably linked to a DNA encoding an interacting protein member. The promoter can be a native promoter, i.e., the promoter found in naturally occurring cells to be responsible for the expression of the interacting protein member in the cells.

Alternatively, the expression cassette can be a chimeric one, i.e., having a heterologous promoter that is not the native promoter responsible for the expression of the interacting protein member in naturally occurring cells. The expression vector may further include an origin of DNA replication for the replication of the vectors in host cells. Preferably, the expression vectors also include a replication origin for the amplification of the vectors in, e.g., *E. coli*, and selection marker(s) for selecting and maintaining only those host cells harboring the expression vectors. Additionally, the expression cassettes preferably also contain inducible elements, which function to control the transcription from the

DNA encoding an interacting protein member. Other regulatory sequences such as transcriptional enhancer sequences and translation regulation sequences (e.g., Shine-Dalgarno sequence) can also be operably included in the expression cassettes.

Termination sequences such as the polyadenylation signals from bovine growth hormone,

5 SV40, lacZ and AcMNPV polyhedral protein genes may also be operably linked to the DNA encoding an interacting protein member in the expression cassettes. An epitope tag coding sequence for detection and/or purification of the expressed protein can also be operably linked to the DNA encoding an interacting protein member such that a fusion protein is expressed. Examples of useful epitope tags include, but are not limited to, 10 influenza virus hemagglutinin (HA), Simian Virus 5 (V5), polyhistidine (6xHis), *c-myc*, lacZ, GST, and the like. Proteins with polyhistidine tags can be easily detected and/or purified with Ni affinity columns, while specific antibodies immunoreactive with many epitope tags are generally commercially available. The expression vectors may also contain components that direct the expressed protein extracellularly or to a particular 15 intracellular compartment. Signal peptides, nuclear localization sequences, endoplasmic reticulum retention signals, mitochondrial localization sequences, myristoylation signals, palmitoylation signals, and transmembrane sequences are example of optional vector components that can determine the destination of expressed proteins. When it is desirable to express two or more interacting protein members in a single host cell, the 20 DNA fragments encoding the interacting protein members may be incorporated into a single vector or different vectors.

The thus constructed expression vectors can be introduced into the host cells by any techniques known in the art, e.g., by direct DNA transformation, microinjection, electroporation, viral infection, lipofection, gene gun, and the like. The expression of the 25 interacting protein members may be transient or stable. The expression vectors can be maintained in host cells in an extrachromosomal state, i.e., as self-replicating plasmids or viruses. Alternatively, the expression vectors can be integrated into chromosomes of the host cells by conventional techniques such as selection of stable cell lines or site-specific recombination. In stable cell lines, at least the expression cassette portion of the 30 expression vector is integrated into a chromosome of the host cells.

The vector construct can be designed to be suitable for expression in various host cells, including but not limited to bacteria, yeast cells, plant cells, insect cells, and mammalian and human cells. Methods for preparing expression vectors for expression in different host cells should be apparent to a skilled artisan.

5 Homologues and fragments of the native interacting protein members can also be easily expressed using the recombinant methods described above. For example, to express a protein fragment, the DNA fragment incorporated into the expression vector can be selected such that it only encodes the protein fragment. Likewise, a specific hybrid protein can be expressed using a recombinant DNA encoding the hybrid protein.

10 Similarly, a homologue protein may be expressed from a DNA sequence encoding the homologue protein. A homologue-encoding DNA sequence may be obtained by manipulating the native protein-encoding sequence using recombinant DNA techniques. For this purpose, random or site-directed mutagenesis can be conducted using techniques generally known in the art. To make protein derivatives, for example, the amino acid

15 sequence of a native interacting protein member may be changed in predetermined manners by site-directed DNA mutagenesis to create or remove consensus sequences for, e.g., phosphorylation by protein kinases, glycosylation, ribosylation, myristolation, palmytoylation, ubiquitination, and the like. Alternatively, non-natural amino acids can be incorporated into an interacting protein member during the synthesis of the protein in

20 recombinant host cells. For example, photoreactive lysine derivatives can be incorporated into an interacting protein member during translation by using a modified lysyl-tRNA. *See, e.g., Wiedmann et al., Nature, 328:830-833 (1989); Musch et al., Cell, 69:343-352 (1992).* Other photoreactive amino acid derivatives can also be incorporated in a similar manner. *See, e.g., High et al., J. Biol. Chem., 368:28745-28751 (1993).*

25 Indeed, the photoreactive amino acid derivatives thus incorporated into an interacting protein member can function to cross-link the protein to its interacting protein partner in a protein complex under predetermined conditions.

In addition, derivatives of the native interacting protein members of the present invention can also be prepared by chemically linking certain moieties to amino acid side

30 chains of the native proteins.

If desired, the homologues and derivatives thus generated can be tested to determine whether they are capable of interacting with their intended partners to form protein complexes. Testing can be conducted by e.g., the yeast two-hybrid system or other methods known in the art for detecting protein-protein interaction.

5 A hybrid protein as described above having survivin or a homologue, derivative, or fragment thereof covalently linked by a peptide bond or a peptide linker to a protein selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT or a homologue, derivative, or fragment thereof, can be expressed recombinantly from a chimeric nucleic acid, e.g., a DNA or mRNA fragment
10 encoding the fusion protein. Accordingly, the present invention also provides a nucleic acid encoding the hybrid protein of the present invention. In addition, an expression vector having incorporated therein a nucleic acid encoding the hybrid protein of the present invention is also provided. The methods for making such chimeric nucleic acids and expression vectors containing them should be apparent to skilled artisans apprised of
15 the present disclosure.

2.4. Protein Microchip

In accordance with another embodiment of the present invention, a protein microchip or microarray is provided having one or more of the protein complexes and/or
20 antibodies selectively immunoreactive with the protein complexes of the present invention. Protein microarrays are becoming increasingly important in both proteomics research and protein-based detection and diagnosis of diseases. The protein microarrays in accordance with this embodiment of the present invention will be useful in a variety of applications including, e.g., large-scale or high-throughput screening for compounds
25 capable of binding to the protein complexes or modulating the interactions between the interacting protein members in the protein complexes.

The protein microarray of the present invention can be prepared in a number of methods known in the art. An example of a suitable method is that disclosed in MacBeath and Schreiber, *Science*, 289:1760-1763 (2000). Essentially, glass microscope
30 slides are treated with an aldehyde-containing silane reagent (SuperAldehyde Substrates purchased from TeleChem International, Cupertino, CA). Nanoliter volumes of protein

samples in a phosphate-buffered saline with 40% glycerol are then spotted onto the treated slides using a high-precision contact-printing robot. After incubation, the slides are immersed in a bovine serum albumin (BSA)-containing buffer to quench the unreacted aldehydes and to form a BSA layer that functions to prevent non-specific protein binding in subsequent applications of the microchip. Alternatively, as disclosed in MacBeath and Schreiber, proteins or protein complexes of the present invention can be attached to a BSA-NHS slide by covalent linkages. BSA-NHS slides are fabricated by first attaching a molecular layer of BSA to the surface of glass slides and then activating the BSA with N,N'-disuccinimidyl carbonate. As a result, the amino groups of the lysine, aspartate, and glutamate residues on the BSA are activated and can form covalent urea or amide linkages with protein samples spotted on the slides. *See* MacBeath and Schreiber, *Science*, 289:1760-1763 (2000).

Another example of a useful method for preparing the protein microchip of the present invention is that disclosed in PCT Publication Nos. WO 00/4389A2 and WO 00/04382, both of which are assigned to Zyomyx and are incorporated herein by reference. First, a substrate or chip base is covered with one or more layers of thin organic film to eliminate any surface defects, insulate proteins from the base materials, and to ensure uniform protein array. Next, a plurality of protein-capturing agents (e.g., antibodies, peptides, etc.) are arrayed and attached to the base that is covered with the thin film. Proteins or protein complexes can then be bound to the capturing agents forming a protein microarray. The protein microchips are kept in flow chambers with an aqueous solution.

The protein microarray of the present invention can also be made by the method disclosed in PCT Publication No. WO 99/36576 assigned to Packard Bioscience Company, which is incorporated herein by reference. For example, a three-dimensional hydrophilic polymer matrix, i.e., a gel, is first dispensed on a solid substrate such as a glass slide. The polymer matrix gel is capable of expanding or contracting and contains a coupling reagent that reacts with amine groups. Thus, proteins and protein complexes can be contacted with the matrix gel in an expanded aqueous and porous state to allow reactions between the amine groups on the protein or protein complexes with the coupling reagents thus immobilizing the proteins and protein complexes on the substrate.

Thereafter, the gel is contracted to embed the attached proteins and protein complexes in the matrix gel.

Alternatively, the proteins and protein complexes of the present invention can be incorporated into a commercially available protein microchip, e.g., the ProteinChip System from CIPHERGEN Biosystems Inc., Palo Alto, CA. The ProteinChip System comprises metal chips having a treated surface, which interact with proteins. Basically, a metal chip surface is coated with a silicon dioxide film. The molecules of interest such as proteins and protein complexes can then be attached covalently to the chip surface via a silane coupling agent.

The protein microchips of the present invention can also be prepared with other methods known in the art, e.g., those disclosed in U.S. Patent Nos. 6,087,102, 6,139,831, 6,087,103; PCT Publication Nos. WO 99/60156, WO 99/39210, WO 00/54046, WO 00/53625, WO 99/51773, WO 99/35289, WO 97/42507, WO 01/01142, WO 00/63694, WO 00/61806, WO 99/61148, WO 99/40434, all of which are incorporated herein by reference.

3. Antibodies

In accordance with another aspect of the present invention, an antibody immunoreactive against a protein complex of the present invention is provided. In one embodiment, the antibody is selectively immunoreactive with a protein complex of the present invention. Specifically, the phrase "selectively immunoreactive with a protein complex" as used herein means that the immunoreactivity of the antibody of the present invention with the protein complex is substantially higher than that with the individual interacting members of the protein complex so that the binding of the antibody to the protein complex is readily distinguishable from the binding of the antibody to the individual interacting member proteins based on the strength of the binding affinities. Preferably, the binding constants differ by a magnitude of at least 2 fold, more preferably at least 5 fold, even more preferably at least 10 fold, and most preferably at least 100 fold. In a specific embodiment, the antibody is not substantially immunoreactive with the interacting protein members of the protein complex.

The antibody of the present invention can be readily prepared using procedures generally known in the art. *See, e.g.,* Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Press, 1988. Typically, the protein complex against which the antibody to be generated will be immunoreactive is used as the antigen for the purpose of producing immune response in a host animal. In one embodiment, the protein complex used consists the native proteins. Preferably, the protein complex includes only the interaction domain(s) of survivin and the interaction domain(s) of one or more proteins selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT. As a result, a greater portion of the total antibodies may be selectively immunoreactive with the protein complexes. The interaction domains can be selected from, e.g., those regions summarized in Table 1. In addition, various techniques known in the art for predicting epitopes may also be employed to design antigenic peptides based on the interacting protein members in a protein complex of the present invention to increase the possibility of producing an antibody selectively immunoreactive with the protein complex. Suitable epitope-prediction computer programs include, e.g., MacVector from International Biotechnologies, Inc. and Protean from DNASTar.

In a specific embodiment, a hybrid protein as described above in Section 2.1 is used as an antigen which has survivin or a homologue, derivative, or fragment thereof covalently linked by a peptide bond or a peptide linker to a protein selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT or a homologue, derivative, or fragment thereof. In a preferred embodiment, the hybrid protein consists of two interacting domains selected from the regions identified in Table 1, or homologues or derivatives thereof, covalently linked together by a peptide bond or a linker molecule.

The antibody of the present invention can be a polyclonal antibody to a protein complex of the present invention. To produce the polyclonal antibody, various animal hosts can be employed, including, e.g., mice, rats, rabbits, goats, guinea pigs, hamsters, etc. A suitable antigen which is a protein complex of the present invention or a derivative thereof as described above can be administered directly to a host animal to illicit immune reactions. Alternatively, it can be administered together with a carrier

such as keyhole limpet hemocyanin (KLH), bovine serum albumin (BSA), ovalbumin, and Tetanus toxoid. Optionally, the antigen is conjugated to a carrier by a coupling agent such as carbodiimide, glutaraldehyde, and MBS. Any conventional adjuvants may be used to boost the immune response of the host animal to the protein complex antigen.

5 Suitable adjuvants known in the art include but are not limited to Complete Freund's Adjuvant (which contains killed mycobacterial cells and mineral oil), incomplete Freund's Adjuvant (which lacks the cellular components), aluminum salts, MF59 from Biocine, monophospholipid, synthetic trehalose dicorynomycolate (TDM) and cell wall skeleton (CWS) both from Corixa Corp., Seattle, WA, non-ionic surfactant vesicles
10 (NISV) from Proteus International PLC, Cheshire, U.K., and saponins. The antigen preparation can be administered to a host animal by subcutaneous, intramuscular, intravenous, intradermal, or intraperitoneal injection, or by injection into a lymphoid organ.

The antibodies of the present invention may also be monoclonal. Such
15 monoclonal antibodies may be developed using any conventional techniques known in the art. For example, the popular hybridoma method disclosed in Kohler and Milstein, *Nature*, 256:495-497 (1975) is now a well-developed technique that can be used in the present invention. See U.S. Patent No. 4,376,110, which is incorporated herein by reference. Essentially, B-lymphocytes producing a polyclonal antibody against a protein
20 complex of the present invention can be fused with myeloma cells to generate a library of hybridoma clones. The hybridoma population is then screened for antigen binding specificity and also for immunoglobulin class (isotype). In this manner, pure hybridoma clones producing specific homogenous antibodies can be selected. See generally, Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Press, 1988.
25 Alternatively, other techniques known in the art may also be used to prepare monoclonal antibodies, which include but are not limited to the EBV hybridoma technique, the human N-cell hybridoma technique, and the trioma technique.

In addition, antibodies selectively immunoreactive with a protein complex of the present invention may also be recombinantly produced. For example, cDNAs prepared
30 by PCR amplification from activated B-lymphocytes or hybridomas may be cloned into an expression vector to form a cDNA library, which is then introduced into a host cell for

recombinant expression. The cDNA encoding a specific desired protein may then be isolated from the library. The isolated cDNA can be introduced into a suitable host cell for the expression of the protein. Thus, recombinant techniques can be used to produce specific native antibodies, hybrid antibodies capable of simultaneous reaction with more than one antigen, chimeric antibodies (e.g., the constant and variable regions are derived from different sources), univalent antibodies that comprise one heavy and light chain pair coupled with the Fc region of a third (heavy) chain, Fab proteins, and the like. See U.S. Patent No. 4,816,567; European Patent Publication No. 0088994; Munro, *Nature*, 312:597 (1984); Morrison, *Science*, 229:1202 (1985); Oi *et al.*, *BioTechniques*, 4:214 (1986); and Wood *et al.*, *Nature*, 314:446-449 (1985), all of which are incorporated herein by reference. Antibody fragments such as Fv fragments, single-chain Fv fragments (scFv), Fab' fragments, and F(ab')₂ fragments can also be recombinantly produced by methods disclosed in, e.g., U.S. Patent No. 4,946,778; Skerra & Plückthun, *Science*, 240:1038-1041(1988); Better *et al.*, *Science*, 240:1041-1043 (1988); and Bird, *et al.*, *Science*, 242:423-426 (1988), all of which are incorporated herein by reference.

In a preferred embodiment, the antibodies provided in accordance with the present invention are partially or fully humanized antibodies. For this purpose, any methods known in the art may be used. For example, partially humanized chimeric antibodies having V regions derived from the tumor-specific mouse monoclonal antibody, but human C regions are disclosed in Morrison and Oi, *Adv. Immunol.*, 44:65-92 (1989). In addition, fully humanized antibodies can be made using transgenic non-human animals. For example, transgenic non-human animals such as transgenic mice can be produced in which endogenous immunoglobulin genes are suppressed or deleted, while heterologous antibodies are encoded entirely by exogenous immunoglobulin genes, preferably human immunoglobulin genes, recombinantly introduced into the genome. See e.g., U.S. Patent Nos. 5,530,101; 5,545,806; 6,075,181; PCT Publication No. WO 94/02602; Green *et al.*, *Nat. Genetics*, 7: 13-21 (1994); and Lonberg *et al.*, *Nature* 368: 856-859 (1994), all of which are incorporated herein by reference. The transgenic non-human host animal may be immunized with suitable antigens such as a protein complex of the present invention or one or more of the interacting protein members thereof to illicit specific immune response thus producing humanized antibodies. In addition, cell lines producing specific

humanized antibodies can also be derived from the immunized transgenic non-human animals. For example, mature B-lymphocytes obtained from a transgenic animal producing humanized antibodies can be fused to myeloma cells and the resulting hybridoma clones may be selected for specific humanized antibodies with desired
5 binding specificities. Alternatively, cDNAs may be extracted from mature B-lymphocytes and used in establishing a library that is subsequently screened for clones encoding humanized antibodies with desired binding specificities.

In yet another embodiment, a bifunctional antibody is provided that has two different antigen binding sites, each being specific to a different interacting protein
10 member in a protein complex of the present invention. The bifunctional antibody may be produced using a variety of methods known in the art. For example, two different monoclonal antibody-producing hybridomas can be fused together. One of the two hybridomas may produce a monoclonal antibody specific against an interacting protein member of a protein complex of the present invention, while the other hybridoma
15 generates a monoclonal antibody immunoreactive with another interacting protein member of the protein complex. The thus formed new hybridoma produces different antibodies including a desired bifunctional antibody, i.e., an antibody immunoreactive with both of the interacting protein members. The bifunctional antibody can be readily purified. *See* Milstein and Cuello, *Nature*, 305:537-540 (1983).

Alternatively, a bifunctional antibody may also be produced using
20 heterobifunctional crosslinkers to chemically link two different monoclonal antibodies, each being immunoreactive with a different interacting protein member of a protein complex. Therefore, the aggregate will bind to two interacting protein members of the protein complex. *See* Staerz *et al*, *Nature*, 314:628-631(1985); Perez *et al*, *Nature*,
25 316:354-356 (1985).

In addition, bifunctional antibodies can also be produced by recombinantly expressing light and heavy chain genes in a hybridoma that itself produces a monoclonal antibody. As a result, a mixture of antibodies including a bifunctional antibody is produced. *See* DeMonte *et al*, *Proc. Natl. Acad. Sci., USA*, 87:2941-2945 (1990); Lenz
30 and Weidle, *Gene*, 87:213-218 (1990).

Preferably, a bifunctional antibody in accordance with the present invention is produced by the method disclosed in U.S. Patent No. 5,582,996, which is incorporated herein by reference. For example, two different Fabs can be provided and mixed together. The first Fab can bind to an interacting protein member of a protein complex, and has a heavy chain constant region having a first complementary domain not naturally present in the Fab but capable of binding a second complementary domain. The second Fab is capable of binding another interacting protein member of the protein complex, and has a heavy chain constant region comprising a second complementary domain not naturally present in the Fab but capable of binding to the first complementary domain. Each of the two complementary domains is capable of stably binding to the other but not to itself. For example, the leucine zipper regions of c-fos and c-jun oncogenes may be used as the first and second complementary domains. As a result, the first and second complementary domains interact with each other to form a leucine zipper thus associating the two different Fabs into a single antibody construct capable of binding to two antigenic sites.

Other suitable methods known in the art for producing bifunctional antibodies may also be used, which include those disclosed in Holliger *et al.*, *Proc. Nat'l Acad. Sci. USA*, 90:6444-6448 (1993); de Kruif *et al.*, *J. Biol. Chem.*, 271:7630-7634 (1996); Coloma and Morrison, *Nat. Biotechnol.*, 15:159-163 (1997); Muller *et al.*, *FEBS Lett.*, 422:259-264 (1998); and Muller *et al.*, *FEBS Lett.*, 432:45-49 (1998), all of which are incorporated herein by reference.

4. Methods of Detecting Protein Complexes

Another aspect of the present invention relates to methods for detecting the protein complexes of the present invention, particularly for determining the concentration of a specific protein complex in a patient sample.

In one embodiment, the concentration of a protein complex having survivin and one or more proteins selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT is determined in cells, tissue, or an organ of a patient. For example, the protein complex can be isolated or purified from a patient sample obtained from cells, tissue, or an organ of the patient and the amount thereof is

determined. As described above, the protein complex can be prepared from cells, tissue or organ samples by coimmunoprecipitation using an antibody immunoreactive with an interacting protein member, a bifunctional antibody that is immunoreactive with two or more interacting protein members of the protein complex, or preferably an antibody
5 selectively immunoreactive with the protein complex. When bifunctional antibodies or antibodies immunoreactive with only free interacting protein members are used, individual interacting protein members not complexed with other proteins may also be isolated along with the protein complex containing such individual proteins. However, they can be readily separated from the protein complex using methods known in the art,
10 e.g., size-based separation methods such as gel filtration, or by subtracting the protein complex from the mixture using an antibody specific against another individual interacting protein member. Additionally, proteins in a sample can be separated in a gel such as polyacrylamide gel and subsequently immunoblotted using an antibody immunoreactive with the protein complex.

15 Alternatively, the concentration of the protein complex can be determined in a sample without separation, isolation or purification. For this purpose, it is preferred that an antibody selectively immunoreactive with the specific protein complex is used in an immunoassay. For example, immunocytochemical methods can be used. Other well known antibody-based techniques can also be used including, e.g., enzyme-linked
20 immunosorbent assay (ELISA), radioimmunoassay (RIA), immunoradiometric assays (IRMA), fluorescent immunoassays, protein A immunoassays, and immunoenzymatic assays (IEMA). *See e.g.*, U.S. Patent Nos. 4,376,110 and 4,486,530, both of which are incorporated herein by reference.

In addition, since a specific protein complex is formed from its interacting protein
25 members, if one of the interacting protein members is at a relatively low concentration in a patient, it may be reasonably expected that the concentration of the protein complex in the patient may also be low. Therefore, the concentration of an individual interacting protein member of a specific protein complex can be determined in a patient sample which can then be used as a reasonably accurate indicator of the concentration of the
30 protein complex in the sample. For this purpose, antibodies against an individual interacting protein member of a specific complex can be used in any one of the methods

described above. In a preferred embodiment, the concentration of each of the interacting protein members of a protein complex is determined in a patient sample and the relative concentration of the protein complex is then deduced.

In addition, the relative protein complex concentration in a patient can also be determined by determining the concentration of the mRNA encoding an interacting protein member of the protein complex. Preferably, each interacting protein member's mRNA concentration in a patient sample is determined. For this purpose, methods for determining mRNA concentration generally known in the art may all be used. Examples of such methods include, e.g., Northern blot assay, dot blot assay, PCR assay (preferably quantitative PCR assay), *in situ* hybridization assay, and the like.

As discussed above, the interactions between survivin and the proteins HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT suggest that these proteins and/or the protein complexes formed by such proteins may be involved in common biological processes and disease pathways. In addition, the interactions between survivin and HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT under physiological conditions may lead to the formation of protein complexes *in vivo* that contain survivin and one or more of the survivin-interacting proteins. The protein complexes are expected to mediate the functions and biological activities of survivin and HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT. For example, survivin and the survivin-interacting proteins may be involved in apoptosis and associated with diseases and disorders such as cancer, viral infection, AIDS, ischemia, stroke, autoimmune diseases, neurodegenerative diseases, inflammatory disorders, sepsis, and osteoporosis. Thus, aberrations in the concentration and/or activity of the protein complexes and/or the proteins such as survivin and the survivin-interacting proteins may result in diseases or disorders such as cancer, viral infection, AIDS, ischemia, stroke, autoimmune diseases, neurodegenerative diseases, inflammatory disorders, sepsis, and osteoporosis. Thus, the aberration in the protein complexes or the individual proteins and the degree of the aberration may be indicators for the diseases or disorders. These aberrations may be used as parameters for classifying and/or staging one of the above-described diseases. In addition, they may also be indicators for patients' response to a drug therapy.

Association between a physiological state (e.g., physiological disorder, predisposition to the disorder, a disease state, response to a drug therapy, or other physiological phenomena or phenotypes) and a specific aberration in a protein complex of the present invention or an individual interacting member thereof can be readily
5 determined by comparative analysis of the protein complex and/or the interacting members thereof in a normal population and an abnormal or affected population. Thus, for example, one can study the concentration, localization and distribution of a particular protein complex, mutations in the interacting protein members of the protein complex, and/or the binding affinity between the interacting protein members in both a normal
10 population and a population affected with a particular physiological disorder described above. The study results can be compared and analyzed by statistical means. Any detected statistically significant difference in the two populations would indicate an association. For example, if the concentration of the protein complex is statistically significantly higher in the affected population than in the normal population, then it can
15 be reasonably concluded that higher concentration of the protein complex is associated with the physiological disorder.

Thus, once an association is established between a particular type of aberration in a particular protein complex of the present invention or in an interacting protein member thereof and a physiological disorder or disease or predisposition to the physiological
20 disorder or disease, then the particular physiological disorder or disease or predisposition to the physiological disorder or disease can be diagnosed or detected by determining whether a patient has the particular aberration.

Accordingly, the present invention also provides a method for diagnosing in a patient a disease or physiological disorder, or a predisposition to the disease or disorder,
25 such as cancer, viral infection, AIDS, ischemia, stroke, autoimmune diseases, neurodegenerative diseases, inflammatory disorders, sepsis, and osteoporosis by determining whether there is any aberration in the patient with respect to a protein complex having a first protein which is survivin interacting with a second protein selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP,
30 OSTP, SLC8A1, A2-CAT. The same protein complex is analyzed in a normal individual and is compared with the results obtained in the patient. In this manner, any protein

complex aberration in the patient can be detected. As used herein, the term “aberration” when used in the context of protein complexes of the present invention means any alterations of a protein complex including increased or decreased concentration of the protein complex in a particular cell or tissue or organ or the total body, altered
5 localization of the protein complex in cellular compartments or in locations of a tissue or organ, changes in binding affinity of an interacting protein member of the protein complex, mutations in an interacting protein member or the gene encoding the protein, and the like. As will be apparent to a skilled artisan, the term “aberration” is used in a relative sense. That is, an aberration is relative to a normal condition.

10 As used herein, the term “diagnosis” means detecting a disease or disorder or determining the stage or degree of a disease or disorder. The term “diagnosis” also encompasses detecting a predisposition to a disease or disorder, determining the therapeutic effect of a drug therapy, or predicting the pattern of response to a drug therapy or xenobiotics. The diagnosis methods of the present invention may be used
15 independently, or in combination with other diagnosing and/or staging methods known in the medical art for a particular disease or disorder.

Thus, in one embodiment, the method of diagnosis is conducted by detecting, in a patient, the concentrations of one or more protein complexes of the present invention using any one of the methods described above, and determining whether the patient has
20 an aberrant concentration of the protein complexes.

The diagnosis may also be based on the determination of the concentrations of one or more interacting protein members (at the protein, cDNA or mRNA level) of a protein complex of the present invention. An aberrant concentration of an interacting protein member may indicate a physiological disorder or a predisposition to a
25 physiological disorder.

In another embodiment, the method of diagnosis comprises determining, in a patient, the cellular localization, or tissue or organ distribution of a protein complex of the present invention and determining whether the patient has an aberrant localization or distribution of the protein complex. For example, immunocytochemical or
30 immunohistochemical assays can be performed on a cell, tissue or organ sample from a patient using an antibody selectively immunoreactive with a protein complex of the

present invention. Antibodies immunoreactive with both an individual interacting protein member and a protein complex containing the protein member may also be used, in which case it is preferred that antibodies immunoreactive with other interacting protein members are also used in the assay. In addition, nucleic acid probes may also be used in
5 *in situ* hybridization assays to detect the localization or distribution of the mRNAs encoding the interacting protein members of a protein complex. Preferably, the mRNA encoding each interacting protein member of a protein complex is detected concurrently.

In yet another embodiment, the method of diagnosis of the present invention comprises detecting any mutations in one or more interacting protein members of a
10 protein complex of the present invention. In particular, it is desirable to determine whether the interacting protein members have any mutations that will lead to, or are associated with, changes in the functional activity of the proteins or changes in their binding affinity to other interacting protein members in forming a protein complex of the present invention. Examples of such mutations include but are not limited to, e.g.,
15 deletions, insertions and rearrangements in the genes encoding the protein members, and nucleotide or amino acid substitutions and the like. In a preferred embodiment, the domains of the interacting protein members that are responsible for the protein-protein interactions, and lead to protein complex formation, are screened to detect any mutations therein. For example, genomic DNA or cDNA encoding an interacting protein member
20 can be prepared from a patient sample, and sequenced. The thus obtained sequence may be compared with known wild-type sequences to identify any mutations. Alternatively, an interacting protein member may be purified from a patient sample and analyzed by protein sequencing or mass spectrometry to detect any amino acid sequence changes. Any methods known in the art for detecting mutations may be used, as will be apparent to
25 skilled artisans apprised of the present disclosure.

In another embodiment, the method of diagnosis includes determining the binding constant of the interacting protein members of one or more protein complexes. For example, the interacting protein members can be obtained from a patient by direct purification or by recombinant expression from genomic DNAs or cDNAs prepared from
30 a patient sample encoding the interacting protein members. Binding constants represent the strength of the protein-protein interaction between the interacting protein members in

a protein complex. Thus, by measuring binding constant, subtle aberration in binding affinity may be detected.

A number of methods known in the art for estimating and determining binding constants in protein-protein interactions are reviewed in Phizicky and Fields, *et al.*, *Microbiol. Rev.*, 59:94-123 (1995), which is incorporated herein by reference. For example, protein affinity chromatography may be used. First, columns are prepared with different concentrations of an interacting protein member which is covalently bound to the columns. Then a preparation of an interacting protein partner is run through the column and washed with buffer. The interacting protein partner bound to the interacting protein member linked to the column is then eluted. Binding constant is then estimated based on the concentrations of the bound protein and the eluted protein. Alternatively, the method of sedimentation through gradients monitors the rate of sedimentation of a mixture of proteins through gradients of glycerol or sucrose. At concentrations above the binding constant, proteins can sediment as a protein complex. Thus, binding constant can be calculated based on the concentrations. Other suitable methods known in the art for estimating binding constant include but are not limited to gel filtration column such as nonequilibrium "small-zone" gel filtration columns (*See e.g.*, Gill *et al.*, *J. Mol. Biol.*, 220:307-324 (1991)), the Hummel-Dreyer method of equilibrium gel filtration (*See e.g.*, Hummel and Dreyer, *Biochim. Biophys. Acta*, 63:530-532 (1962)) and large-zone equilibrium gel filtration (*See e.g.*, Gilbert and Kellett, *J. Biol. Chem.*, 246:6079-6086 (1971)), sedimentation equilibrium (*See e.g.*, Rivas and Minton, *Trends Biochem.*, 18:284-287 (1993)), fluorescence methods such as fluorescence spectrum (*See e.g.*, Otto-Bruc *et al.*, *Biochemistry*, 32:8632-8645 (1993)) and fluorescence polarization or anisotropy with tagged molecules (*See e.g.*, Weiel and Hershey, *Biochemistry*, 20:5859-5865 (1981)), solution equilibrium measured with immobilized binding protein (*See e.g.*, Nelson and Long, *Biochemistry*, 30:2384-2390 (1991)), and surface plasmon resonance (*See e.g.*, Panayotou *et al.*, *Mol. Cell. Biol.*, 13:3567-3576 (1993)).

In another embodiment, the diagnosis method of the present invention comprises detecting protein-protein interactions in functional assay systems such as the yeast two-hybrid system. Accordingly, to determine the protein-protein interaction between two interacting protein members that normally form a protein complex in normal individuals,

cDNAs encoding the interacting protein members can be isolated from a patient to be diagnosed. The thus cloned cDNAs or fragments thereof can be subcloned into vectors for use in yeast two-hybrid system. Preferably a reverse yeast two-hybrid system is used such that failure of interaction between the proteins may be positively detected. The use
5 of yeast two-hybrid system or other systems for detecting protein-protein interactions is known in the art and is described below in Section 5.3.1.

A kit may be used for conducting the diagnosis methods of the present invention. Typically, the kit should contain, in a carrier or compartmentalized container, reagents useful in any of the above-described embodiments of the diagnosis method. The carrier
10 can be a container or support, in the form of, e.g., bag, box, tube, rack, and is optionally compartmentalized. The carrier may define an enclosed confinement for safety purposes during shipment and storage. In one embodiment, the kit includes an antibody selectively immunoreactive with a protein complex of the present invention. In addition, antibodies against individual interacting protein members of the protein complexes may also be
15 included. The antibodies may be labeled with a detectable marker such as radioactive isotopes, or enzymatic or fluorescence markers. Alternatively secondary antibodies such as labeled anti-IgG and the like may be included for detection purposes. Optionally, the kit can include one or more of the protein complexes of the present invention prepared or purified from a normal individual or an individual afflicted with a physiological disorder associated with an aberration in the protein complexes or an interacting protein member thereof. In addition, the kit may further include one or more of the interacting protein members of the protein complexes of the present invention prepared or purified from a normal individual or an individual afflicted with a physiological disorder associated with an aberration in the protein complexes or an interacting protein member thereof. Suitable
20 oligonucleotide primers useful in the amplification of the genes or cDNAs for the interacting protein members may also be provided in the kit. In particular, in a preferred embodiment, the kit includes a first oligonucleotide selectively hybridizable to the mRNA or cDNA encoding survivin and a second oligonucleotide selectively hybridizable to the mRNA or cDNA encoding a protein selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT. Additional
25 30 oligonucleotides hybridizing to a region of the gene encoding survivin and/or a region of

the gene(s) encoding one or more survivin-interacting proteins, as identified in the present invention, may also be included. Such oligonucleotides may be used as PCR primers for, e.g., quantitative PCR amplification of mRNAs encoding survivin and an interacting partner thereof, or as hybridizing probes for detecting the mRNAs. The
5 oligonucleotides may have a length of from about 8 nucleotides to about 100 nucleotides, preferably from about 12 to about 50 nucleotides, and more preferably from about 15 to about 30 nucleotides. In addition, the kit may also contain oligonucleotides that can be used as hybridization probes for detecting the cDNAs or mRNAs encoding the interacting protein members. Preferably, instructions for using the kit or reagents contained therein
10 are also included in the kit.

5. Use of Protein Complexes or Interacting Protein Members Thereof in Screening Assays for Modulators

The protein complexes of the present invention and survivin and survivin-
15 interacting proteins such as HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT can also be used in screening assays to identify modulators of the protein complexes, survivin, and/or the survivin-interacting proteins. In addition, homologues, derivatives and fragments of survivin and homologues, derivatives and fragments of the survivin-interacting proteins may also be used in such screening assays.
20 As used herein, the term “modulator” encompasses any compounds that can cause any forms of alteration of the biological activities or functions of the proteins or protein complexes, including, e.g., enhancing or reducing their biological activities, increasing or decreasing their stability, altering their affinity or specificity to certain other biological molecules, etc. In addition, the term “modulator” as used herein also includes any
25 compounds that simply bind survivin, survivin-interacting proteins, and/or the proteins complexes of the present invention. For example, a modulator can be an “interaction antagonist” capable of interfering with or disrupting or dissociating protein-protein interaction between survivin or a homologue, fragment or derivative thereof and one or more proteins selected from the group consisting of HDLC1, beta-actin, DNA helicase II,
30 COPP, OSTP, SLC8A1, A2-CAT or a homologue, fragment or derivative thereof. A modulator can also be an “interaction agonist” that initiates or strengthens the interaction

between the protein members of a protein complex of the present invention, or homologues, fragments or derivatives thereof.

Accordingly, the present invention provides screening methods for selecting modulators of survivin, or a mutant form thereof, a survivin-interacting protein selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT, or a mutant form thereof, or a protein complex formed between survivin, or a mutant form thereof, and one or more of the survivin-interacting proteins, or a mutant forms thereof. Screening methods are also provided for selecting modulators of survivin homologues, derivatives or fragments, or homologues, derivatives or fragments of a survivin-interacting protein, or a protein complex formed between a survivin homologue, derivative or fragment and a homologue or derivative or fragment of a survivin-interacting protein, or proteins.

The modulators selected in accordance with the screening methods of the present invention can be effective in modulating the functions or activities of survivin, a survivin-interacting protein, or the protein complexes of the present invention. For example, compounds capable of binding to the protein complexes may be capable of modulating the functions of the protein complexes. Additionally, compounds that interfere with, weaken, dissociate or disrupt, or alternatively, initiate, facilitate or stabilize the protein-protein interaction between the interacting protein members of the protein complexes can also be effective in modulating the functions or activities of the protein complexes. Thus, the compounds identified in the screening methods of the present invention can be made into therapeutically or prophylactically effective drugs for preventing or ameliorating diseases, disorders or symptoms caused by or associated with aberrations in the protein complexes or survivin or the survivin-interacting proteins of the present invention.

Alternatively, they may be used as leads to aid the design and identification of therapeutically or prophylactically effective compounds for diseases, disorders or symptoms caused by or associated with aberrations in the protein complexes or survivin or the survivin-interacting proteins of the present invention. The protein complexes and/or interacting protein members thereof in accordance with the present invention can be used in any of a variety of drug screening techniques. Drug screening can be performed as described herein or using well-known techniques, such as those described

in U.S. Patent Nos. 5,800,998 and 5,891,628, both of which are incorporated herein by reference.

5.1. Test Compounds

Any test compounds may be screened in the screening assays of the present invention to select modulators of survivin, a survivin-containing protein complex and/or a survivin-interacting protein of the present invention. By the term “selecting” or “select” compounds it is intended to encompass both (a) choosing compounds from a group previously unknown to be modulators of survivin, a survivin-containing protein complex and/or a survivin-interacting protein of the present invention, and (b) testing compounds that are known to be capable of binding, or modulating the functions and activities of, survivin, a survivin-containing protein complex and/or a survivin-interacting protein of the present invention. Both types of compounds are generally referred to herein as “test compounds.” The test compounds may include, by way of example, proteins (e.g., antibodies, small peptides, artificial or natural proteins), nucleic acids, and derivatives, mimetics and analogs thereof, and small organic molecules having a molecular weight of no greater than 10,000 daltons, more preferably less than 5,000 daltons. Preferably, the test compounds are provided in library formats known in the art, e.g., in chemically synthesized libraries, recombinantly expressed libraries (e.g., phage display libraries), and *in vitro* translation-based libraries (e.g., ribosome display libraries).

For example, the screening assays of the present invention can be used in the antibody production processes described in Section 3 to select antibodies with desirable specificities. Various forms antibodies or derivatives thereof may be screened, including but not limited to, polyclonal antibodies, monoclonal antibodies, bifunctional antibodies, chimeric antibodies, single chain antibodies, antibody fragments such as Fv fragments, single-chain Fv fragments (scFv), Fab' fragments, and F(ab')₂ fragments, and various modified forms of antibodies such as catalytic antibodies, and antibodies conjugated to toxins or drugs, and the like. The antibodies can be of any types such as IgG, IgE, IgA, or IgM. Humanized antibodies are particularly preferred. Preferably, the various antibodies and antibody fragments may be provided in libraries to allow large-scale high throughput screening. For example, expression libraries expressing antibodies or

antibody fragments may be constructed by a method disclosed, e.g., in Huse *et al.*, *Science*, 246:1275-1281 (1989), which is incorporated herein by reference. Single-chain Fv (scFv) antibodies are of particular interest in diagnostic and therapeutic applications. Methods for providing antibody libraries are also provided in U.S. Patent Nos. 6,096,551; 5,844,093; 5,837,460; 5,789,208; and 5,667,988, all of which are incorporated herein by reference.

Peptidic test compounds may be peptides having L-amino acids and/or D-amino acids, phosphopeptides, and other types of peptides. The screened peptides can be of any size, but preferably have less than about 50 amino acids. Smaller peptides are easier to deliver into a patient's body. Various forms of modified peptides may also be screened. Like antibodies, peptides can also be provided in, e.g., combinatorial libraries. *See generally*, Gallop *et al.*, *J. Med. Chem.*, 37:1233-1251 (1994). Methods for making random peptide libraries are disclosed in, e.g., Devlin *et al.*, *Science*, 249:404-406 (1990). Other suitable methods for constructing peptide libraries and screening peptides therefrom are disclosed in, e.g., Scott and Smith, *Science*, 249:386-390 (1990); Moran *et al.*, *J. Am. Chem. Soc.*, 117:10787-10788 (1995) (a library of electronically tagged synthetic peptides); Stachelhaus *et al.*, *Science*, 269:69-72 (1995); U.S. Patent Nos. 6,156,511; 6,107,059; 6,015,561; 5,750,344; 5,834,318; 5,750,344, all of which are incorporated herein by reference. For example, random-sequence peptide phage display libraries may be generated by cloning synthetic oligonucleotides into the gene III or gene VIII of an *E. coli*. filamentous phage. The thus generated phage can propagate in *E. coli*. and express peptides encoded by the oligonucleotides as fusion proteins on the surface of the phage. Scott and Smith, *Science*, 249:368-390 (1990). Alternatively, the "peptides on plasmids" method may also be used to form peptide libraries. In this method, random peptides may be fused to the C-terminus of the *E. coli*. Lac repressor by recombinant technologies and expressed from a plasmid that also contains Lac repressor-binding sites. As a result, the peptide fusions bind to the same plasmid that encodes them.

Small organic or inorganic non-peptide non-nucleotide compounds are preferred test compounds for the screening assays of the present invention. They too can be provided in a library format. *See generally*, Gordan *et al.* *J. Med. Chem.*, 37:1385-1401 (1994). For example, benzodiazepine libraries are provided in Bunin and Ellman, *J. Am.*

Chem. Soc., 114:10997-10998 (1992), which is incorporated herein by reference. A method for constructing and screening peptoid libraries are disclosed in Simon *et al.*, *Proc. Natl. Acad. Sci. USA*, 89:9367-9371 (1992). Methods for the biosynthesis of novel polyketides in a library format are described in McDaniel *et al.*, *Science*, 262:1546-1550 (1993) and Kao *et al.*, *Science*, 265:509-512 (1994). Various libraries of small organic molecules and methods of construction thereof are disclosed in U.S. Patent Nos. 6,162,926 (multiply-substituted fullerene derivatives); 6,093,798 (hydroxamic acid derivatives); 5,962,337 (combinatorial 1,4-benzodiazepin-2, 5-dione library); 5,877,278 (Synthesis of N-substituted oligomers); 5,866,341 (compositions and methods for screening drug libraries); 5,792,821 (polymerizable cyclodextrin derivatives); 5,766,963 (hydroxypropylamine library); and 5,698,685 (morpholino-subunit combinatorial library), all of which are incorporated herein by reference.

Other compounds such as oligonucleotides and peptide nucleic acids (PNA), and analogs and derivatives thereof may also be screened to identify clinically useful compounds. Combinatorial libraries of oligonucleotides are also known in the art. *See* Gold *et al.*, *J. Biol. Chem.*, 270:13581-13584 (1995).

5.2. *In vitro* Screening Assays

The test compounds may be screened in an *in vitro* assay to identify compounds capable of binding the protein complexes or interacting protein members thereof in accordance with the present invention. For this purpose, a test compound is contacted with a protein complex or an interacting protein member thereof under conditions and for a time sufficient to allow specific interaction between the test compound and the target components to occur and thus binding of the compound to the target forming a complex. Subsequently, the binding event is detected.

Various screening techniques known in the art may be used in the present invention. The protein complexes and the interacting protein members thereof may be prepared by any suitable methods, e.g., by recombinant expression and purification. The protein complexes and/or interacting protein members thereof (both are referred to as “target” hereinafter in this section) may be free in solution. A test compound may be mixed with a target forming a liquid mixture. The compound may be labeled with a

detectable marker. Upon mixing under suitable conditions, the binding complex having the compound and the target may be co-immunoprecipitated and washed. The compound in the precipitated complex may be detected based on the marker on the compound.

In a preferred embodiment, the target is immobilized on a solid support or on a cell surface. Preferably, the target can be arrayed into a protein microchip in a method described in Section 2.3. For example, a target may be immobilized directly onto a microchip substrate such as glass slides or onto a multi-well plates using non-neutralizing antibodies, i.e., antibodies capable of binding to the target but do not substantially affect its biological activities. To affect the screening, test compounds can be contacted with the immobilized target to allow binding to occur to form complexes under standard binding assay conditions. Either the targets or test compounds are labeled with a detectable marker using well-known labeling techniques. For example, U.S. Patent No. 5,741,713 discloses combinatorial libraries of biochemical compounds labeled with NMR active isotopes. To identify binding compounds, one may measure the formation of the target-test compound complexes or kinetics for the formation thereof. When combinatorial libraries of organic non-peptide non-nucleic acid compound are screened, it is preferred that labeled or encoded (or "tagged") combinatorial libraries are used to allow rapid decoding of lead structures. This is especially important because, unlike biological libraries, individual compounds found in chemical libraries cannot be amplified by self-amplification. Tagged combinatorial libraries are provided in, e.g., Borchardt and Still, *J. Am. Chem. Soc.*, 116:373-374 (1994) and Moran *et al.*, *J. Am. Chem. Soc.*, 117:10787-10788 (1995), both of which are incorporated herein by reference.

Alternatively, the test compounds can be immobilized on a solid support, e.g., forming a microarray of test compounds. The target protein or protein complex is then contacted with the test compounds. The target may be labeled with any suitable detection marker. For example, the target may be labeled with radioactive isotopes or fluorescence marker before binding reaction occurs. Alternatively, after the binding reactions, antibodies that are immunoreactive with the target and are labeled with radioactive materials, fluorescence markers, enzymes, or labeled secondary anti-Ig antibodies may be used to detect any bound target thus identifying the binding compound. One example of

this embodiment is the protein probing method. That is, the target provided in accordance with the present invention is used as a probe to screen expression libraries of proteins or random peptides. The expression libraries can be phage display libraries, *in vitro* translation-based libraries, or ordinary expression cDNA libraries. The libraries
5 may be immobilized on a solid support such as nitrocellulose filters. *See e.g.*, Sikela and Hahn, *Proc. Natl. Acad. Sci. USA*, 84:3038-3042 (1987). The probe may be labeled with a radioactive isotope or a fluorescence marker. Alternatively, the probe can be biotinylated and detected with a streptavidin-alkaline phosphatase conjugate. More conveniently, the bound probe may be detected with an antibody.

10 In yet another embodiment, a known ligand capable of binding to the target can be used in competitive binding assays. Complexes between the known ligand and the target can be formed and then contacted with test compounds. The ability of a test compound to interfere with the interaction between the target and the known ligand is measured. One exemplary ligand is an antibody capable of specifically binding the
15 target. Particularly, such an antibody is especially useful for identifying peptides that share one or more antigenic determinants of the target protein complex or interacting protein members thereof.

In a specific embodiment, a protein complex used in the screening assay includes a hybrid protein as described in Section 2.1, which is formed by fusion of two interacting
20 protein members or fragments or interaction domains thereof. The hybrid protein may also be designed such that it contains a detectable epitope tag fused thereto. Suitable examples of such epitope tags include sequences derived from, e.g., influenza virus hemagglutinin (HA), Simian Virus 5 (V5), polyhistidine (6xHis), *c-myc*, lacZ, GST, and the like.

25 Test compounds may be also screened in an *in vitro* assay to identify compounds capable of dissociating the protein complexes identified in accordance with the present invention. Thus, for example, a survivin-containing protein complex can be contacted with a test compound and the protein complex can be detected. Conversely, test compounds may also be screened to identify compounds capable of enhancing the
30 interaction between survivin and a survivin-interacting protein or stabilizing the protein complex formed by the two or more proteins.

The assay can be conducted in similar manners as the binding assays described above. For example, the presence or absence of a particular protein complex can be detected by an antibody selectively immunoreactive with the protein complex. Thus, after incubation of the protein complex with a test compound, an immunoprecipitation assay can be conducted with the antibody. If the test compound disrupts the protein complex, then the amount of immunoprecipitated protein complex in this assay will be significantly less than that in a control assay in which the same protein complex is not contacted with the test compound. Similarly, two proteins the interaction between which is to be enhanced may be incubated together with a test compound. Thereafter, a protein complex may be detected by the selectively immunoreactive antibody. The amount of protein complex may be compared to that formed in the absence of the test compound. Various other detection methods may be suitable in the dissociation assay, as will be apparent to skilled artisan apprised of the present disclosure.

5.3. *In vivo* Screening Assays

Test compounds can also be screened in any *in vivo* assays to select modulators of the protein complexes or interacting protein members thereof in accordance with the present invention. For example, any *in vivo* assays known in the art to be useful in identifying compounds capable of strengthening or interfering with the stability of the protein complexes of the present invention may be used.

5.3.1. Two-Hybrid Assays

In a preferred embodiment, one of the yeast two-hybrid systems or their analogous or derivative forms is used. Examples of suitable two-hybrid systems known in the art include, but are not limited to, those disclosed in U.S. Patent Nos. 5,283,173; 5,525,490; 5,585,245; 5,637,463; 5,695,941; 5,733,726; 5,776,689; 5,885,779; 5,905,025; 6,037,136; 6,057,101; 6,114,111; and Bartel and Fields, eds., *The Yeast Two-Hybrid System*, Oxford University Press, New York, NY, 1997, all of which are incorporated herein by reference.

Typically, in a classic transcription-based two-hybrid assay, two chimeric genes are prepared encoding two fusion proteins: one contains a transcription activation domain

204766-1266007
fused to an interacting protein member of a protein complex of the present invention or
an interaction domain or fragment of the interacting protein member, while the other
fusion protein includes a DNA binding domain fused to another interacting protein
member of the protein complex or a fragment or interaction domain thereof. For the
5 purpose of convenience, the two interacting protein members, fragments or interaction
domains thereof are referred to as "bait fusion protein" and "prey fusion protein,"
respectively. The chimeric genes encoding the fusion proteins are termed "bait chimeric
gene" and "prey chimeric gene," respectively. Typically, a "bait vector" and a "prey
vector" are provided for the expression of a bait chimeric gene and a prey chimeric gene,
10 respectively.

5.3.1.1. Vectors

Many types of vectors can be used in a transcription-based two-hybrid assay.
Methods for the construction of bait vectors and prey vectors should be apparent to
15 skilled artisans in the art apprised of the present disclosure. *See generally, Current
Protocols in Molecular Biology*, Vol. 2, Ed. Ausubel, *et al.*, Greene Publish. Assoc. &
Wiley Interscience, Ch. 13, 1988; Glover, *DNA Cloning*, Vol. II, IRL Press, Wash., D.C.,
Ch. 3, 1986; Bitter, *et al.*, in *Methods in Enzymology* 153:516-544 (1987); *The Molecular
Biology of the Yeast Saccharomyces*, Eds. Strathern *et al.*, Cold Spring Harbor Press,
20 Vols. I and II, 1982; and Rothstein in *DNA Cloning: A Practical Approach*, Vol. 11, Ed.
DM Glover, IRL Press, Wash., D.C., 1986.

Generally, the bait and prey vectors include an expression cassette having a
promoter operably linked to a chimeric gene for the transcription of the chimeric gene.
The vectors may also include an origin of DNA replication for the replication of the
25 vectors in host cells and a replication origin for the amplification of the vectors in, e.g., *E.
coli*, and selection marker(s) for selecting and maintaining only those host cells harboring
the vectors. Additionally, the expression cassette preferably also contains inducible
elements, which function to control the expression of a chimeric gene. Making the
expression of the chimeric genes inducible and controllable is especially important in the
30 event that the fusion proteins or components thereof are toxic to the host cells. Other
regulatory sequences such as transcriptional enhancer sequences and translation

regulation sequences (e.g., Shine-Dalgarno sequence) can also be included in the expression cassette. Termination sequences such as the bovine growth hormone, SV40, lacZ and AcMNPV polyhedral polyadenylation signals may also be operably linked to a chimeric gene in the expression cassette. An epitope tag coding sequence for detection and/or purification of the fusion proteins can also be operably linked to the chimeric gene in the expression cassette. Examples of useful epitope tags include, but are not limited to, influenza virus hemagglutinin (HA), Simian Virus 5 (V5), polyhistidine (6xHis), *c-myc*, lacZ, GST, and the like. Proteins with polyhistidine tags can be easily detected and/or purified with Ni affinity columns, while specific antibodies to many epitope tags are generally commercially available. The vectors can be introduced into the host cells by any techniques known in the art, e.g., by direct DNA transformation, microinjection, electroporation, viral infection, lipofection, gene gun, and the like. The bait and prey vectors can be maintained in host cells in an extrachromosomal state, i.e., as self-replicating plasmids or viruses. Alternatively, one or both vectors can be integrated into chromosomes of the host cells by conventional techniques such as selection of stable cell lines or site-specific recombination.

The *in vivo* assays of the present invention can be conducted in many different host cells, including but not limited to bacteria, yeast cells, plant cells, insect cells, and mammalian cells. A skilled artisan will recognize that the designs of the vectors can vary with the host cells used. In one embodiment, the assay is conducted in prokaryotic cells such as *Escherichia coli*, *Salmonella*, *Klebsiella*, *Pseudomonas*, *Caulobacter*, and *Rhizobium*. Suitable origins of replication for the expression vectors useful in this embodiment of the present invention include, e.g., the ColE1, pSC101, and M13 origins of replication. Examples of suitable promoters include, for example, the T7 promoter, the lacZ promoter, and the like. In addition, inducible promoters are also useful in modulating the expression of the chimeric genes. For example, the lac operon from bacteriophage lambda plac5 is well known in the art and is inducible by the addition of IPTG to the growth medium. Other known inducible promoters useful in a bacteria expression system include pL of bacteriophage λ , the trp promoter, and hybrid promoters such as the tac promoter, and the like.

In addition, selection marker sequences for selecting and maintaining only those prokaryotic cells expressing the desirable fusion proteins should also be incorporated into the expression vectors. Numerous selection markers including auxotrophic markers and antibiotic resistance markers are known in the art and can all be useful for purposes of this invention. For example, the *bla* gene, which confers ampicillin resistance, is the most commonly used selection marker in prokaryotic expression vectors. Other suitable markers include genes that confer neomycin, kanamycin, or hygromycin resistance to the host cells. In fact, many vectors are commercially available from vendors such as Invitrogen Corp. of Carlsbad, CA, Clontech Corp. of Palo Alto, CA, and Stratagene Corp. of La Jolla, CA, and Promega Corp. of Madison, WI. These commercially available vectors, e.g., pBR322, pSPORT, pBluescriptIISK, pcDNAI, and pcDNAII all have a multiple cloning site into which the chimeric genes of the present invention can be conveniently inserted using conventional recombinant techniques. The constructed expression vectors can be introduced into host cells by various transformation or transfection techniques generally known in the art.

In another embodiment, mammalian cells are used as host cells for the expression of the fusion proteins and detection of protein-protein interactions. For this purpose, virtually any mammalian cells can be used including normal tissue cells, stable cell lines, and transformed tumor cells. Conveniently, mammalian cell lines such as CHO cells, Jurkat T cells, NIH 3T3 cells, HEK-293 cells, CV-1 cells, COS-1 cells, HeLa cells, VERO cells, MDCK cells, WI38 cells, and the like are used. Mammalian expression vectors are well known in the art and many are commercially available. Examples of suitable promoters for the transcription of the chimeric genes in mammalian cells include viral transcription promoters derived from adenovirus, simian virus 40 (SV40) (e.g., the early and late promoters of SV40), Rous sarcoma virus (RSV), and cytomegalovirus (CMV) (e.g., CMV immediate-early promoter), human immunodeficiency virus (HIV) (e.g., long terminal repeat (LTR)), vaccinia virus (e.g., 7.5K promoter), and herpes simplex virus (HSV) (e.g., thymidine kinase promoter). Inducible promoters can also be used. Suitable inducible promoters include, for example, the tetracycline responsive element (TRE) (*See Gossen et al., Proc. Natl. Acad. Sci. USA, 89:5547-5551 (1992)*), metallothionein IIA promoter, ecdysone-responsive promoter, and heat shock promoters.

Suitable origins of replication for the replication and maintenance of the expression vectors in mammalian cells include, e.g., the Epstein Barr origin of replication in the presence of the Epstein Barr nuclear antigen (*see* Sugden *et al.*, *Mole. Cell. Biol.*, 5:410-413 (1985)) and the SV40 origin of replication in the presence of the SV40 T antigen (which is present in COS-1 and COS-7 cells) (*see* Margolskee *et al.*, *Mole. Cell. Biol.*, 8:2837 (1988)). Suitable selection markers include, but are not limited to, genes conferring resistance to neomycin, hygromycin, zeocin, and the like. Many commercially available mammalian expression vectors may be useful for the present invention, including, e.g., pCEP4, pcDNA1, pIND, pSecTag2, pVAX1, pcDNA3.1, and pBI-EGFP, and pDisplay. The vectors can be introduced into mammalian cells using any known techniques such as calcium phosphate precipitation, lipofection, electroporation, and the like. The bait vector and prey vector can be co-transformed into the same cell or, alternatively, introduced into two different cells which are subsequently fused together by cell fusion or other suitable techniques.

Viral expression vectors, which permit introduction of recombinant genes into cells by viral infection, can also be used for the expression of the fusion proteins. Viral expression vectors generally known in the art include viral vectors based on adenovirus, bovine papilloma virus, murine stem cell virus (MSCV), MFG virus, and retrovirus. *See* Sarver, *et al.*, *Mol. Cell. Biol.*, 1: 486 (1981); Logan & Shenk, *Proc. Natl. Acad. Sci. USA*, 81:3655-3659 (1984); Mackett, *et al.*, *Proc. Natl. Acad. Sci. USA*, 79:7415-7419 (1982); Mackett, *et al.*, *J. Virol.*, 49:857-864 (1984); Panicali, *et al.*, *Proc. Natl. Acad. Sci. USA*, 79:4927-4931 (1982); Cone & Mulligan, *Proc. Natl. Acad. Sci. USA*, 81:6349-6353 (1984); Mann *et al.*, *Cell*, 33:153-159 (1993); Pear *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:8392-8396 (1993); Kitamura *et al.*, *Proc. Natl. Acad. Sci. USA*, 92:9146-9150 (1995); Kinsella *et al.*, *Human Gene Therapy*, 7:1405-1413 (1996); Hofmann *et al.*, *Proc. Natl. Acad. Sci. USA*, 93:5185-5190 (1996); Choate *et al.*, *Human Gene Therapy*, 7:2247 (1996); WO 94/19478; Hawley *et al.*, *Gene Therapy*, 1:136 (1994) and Rivere *et al.*, *Genetics*, 92:6733 (1995), all of which are incorporated by reference.

Generally, to construct a viral vector, a chimeric gene according to the present invention can be operably linked to a suitable promoter. The promoter-chimeric gene construct is then inserted into a non-essential region of the viral vector, typically a

modified viral genome. This results in a viable recombinant virus capable of expressing the fusion protein encoded by the chimeric gene in infected host cells. Once in the host cell, the recombinant virus typically is integrated into the genome of the host cell.

However, recombinant bovine papilloma viruses typically replicate and remain as extrachromosomal elements.

In another embodiment, the detection assays of the present invention are conducted in plant cell systems. Methods for expressing exogenous proteins in plant cells are well known in the art. *See generally*, Weissbach & Weissbach, *Methods for Plant Molecular Biology*, Academic Press, NY, 1988; Grierson & Corey, *Plant Molecular Biology*, 2d Ed., Blackie, London, 1988. Recombinant virus expression vectors based on, e.g., cauliflower mosaic virus (CaMV) or tobacco mosaic virus (TMV) can all be used. Alternatively, recombinant plasmid expression vectors such as Ti plasmid vectors and Ri plasmid vectors are also useful. The chimeric genes encoding the fusion proteins of the present invention can be conveniently cloned into the expression vectors and placed under control of a viral promoter such as the 35S RNA and 19S RNA promoters of CaMV or the coat protein promoter of TMV, or of a plant promoter, e.g., the promoter of the small subunit of RUBISCO and heat shock promoters (e.g., soybean hsp17.5-E or hsp17.3-B promoters).

In addition, the *in vivo* assay of the present invention can also be conducted in insect cells, e.g., *Spodoptera frugiperda* cells, using a baculovirus expression system. Expression vectors and host cells useful in this system are well known in the art and are generally available from various commercial vendors. For example, the chimeric genes of the present invention can be conveniently cloned into a non-essential region (e.g., the polyhedrin gene) of an *Autographa californica* nuclear polyhedrosis virus (AcNPV) vector and placed under control of an AcNPV promoter (e.g., the polyhedrin promoter). The non-occluded recombinant viruses thus generated can be used to infect host cells such as *Spodoptera frugiperda* cells in which the chimeric genes are expressed. *See* U.S. Patent No. 4,215,051.

In a preferred embodiment of the present invention, the fusion proteins are expressed in a yeast expression system using yeasts such as *Saccharomyces cerevisiae*, *Hansenula polymorpha*, *Pichia pastoris*, and *Schizosaccharomyces pombe* as host cells.

The expression of recombinant proteins in yeasts is a well-developed field, and the techniques useful in this respect are disclosed in detail in *The Molecular Biology of the Yeast Saccharomyces*, Eds. Strathern *et al.*, Vols. I and II, Cold Spring Harbor Press, 1982; Ausubel *et al.*, *Current Protocols in Molecular Biology*, New York, Wiley, 1994; and Guthrie and Fink, *Guide to Yeast Genetics and Molecular Biology*, in *Methods in Enzymology*, Vol. 194, 1991, all of which are incorporated herein by reference. Sudbery, *Curr. Opin. Biotech.*, 7:517-524 (1996) reviews the success in the art in expressing recombinant proteins in various yeast species; the entire content and references cited therein are incorporated herein by reference. In addition, Bartel and Fields, eds., *The Yeast Two-Hybrid System*, Oxford University Press, New York, NY, 1997 contains extensive discussions of recombinant expression of fusion proteins in yeasts in the context of various yeast two-hybrid systems, and cites numerous relevant references. These and other methods known in the art can all be used for purposes of the present invention. The application of such methods to the present invention should be apparent to a skilled artisan apprised of the present disclosure.

Generally, each of the two chimeric genes is included in a separate expression vector (bait vector and prey vector). Both vectors can be co-transformed into a single yeast host cell. As will be apparent to a skilled artisan, it is also possible to express both chimeric genes from a single vector. In a preferred embodiment, the bait vector and prey vector are introduced into two haploid yeast cells of opposite mating types, e.g., a-type and α -type, respectively. The two haploid cells can be mated at a desired time to form a diploid cell expressing both chimeric genes.

Generally, the bait and prey vectors for recombinant expression in yeast include a yeast replication origin such as the 2μ origin or the *ARSH4* sequence for the replication and maintenance of the vectors in yeast cells. Preferably, the vectors also have a bacteria origin of replication (e.g., ColE1) and a bacteria selection marker (e.g., amp^R marker, i.e., *bla* gene). Optionally, the *CEN6* centromeric sequence is included to control the replication of the vectors in yeast cells. Any constitutive or inducible promoters capable of driving gene transcription in yeast cells may be employed to control the expression of the chimeric genes. Such promoters are operably linked to the chimeric genes. Examples of suitable constitutive promoters include but are not limited to the yeast *ADHI*, *PGKI*,

TEF2 , GPD1 , HIS3, and CYC1 promoters. Example of suitable inducible promoters include but are not limited to the yeast GAL1 (inducible by galactose), CUP1 (inducible by Cu⁺⁺), and FUS1 (inducible by pheromone) promoters; the AOX/MOX promoter from *H. polymorpha* and *P. Pastoris* (repressed by glucose or ethanol and induced by methanol); chimeric promoters such as those that contain LexA operators (inducible by LexA-containing transcription factors); and the like. Inducible promoters are preferred when the fusion proteins encoded by the chimeric genes are toxic to the host cells. If it is desirable, certain transcription repressing sequences such as the upstream repressing sequence (URS) from SPO13 promoter can be operably linked to the promoter sequence, e.g., to the 5' end of the promoter region. Such upstream repressing sequences function to fine-tune the expression level of the chimeric genes.

Preferably, a transcriptional termination signal is operably linked to the chimeric genes in the vectors. Generally, transcriptional termination signal sequences derived from, e.g., the *CYC1* and *ADH1* genes can be used.

Additionally, it is preferred that the bait vector and prey vector contain one or more selectable markers for the selection and maintenance of only those yeast cells that harbor one or both chimeric genes. Any selectable markers known in the art can be used for purposes of this invention so long as yeast cells expressing the chimeric gene(s) can be positively identified or negatively selected. Examples of markers that can be positively identified are those based on color assays, including the lacZ gene (which encodes β -galactosidase), the firefly luciferase gene, secreted alkaline phosphatase, horseradish peroxidase, the blue fluorescent protein (BFP), and the green fluorescent protein (GFP) gene (*see* Cubitt *et al.*, *Trends Biochem. Sci.*, 20:448-455 (1995)). Other markers allowing detection by fluorescence, chemiluminescence, UV absorption, infrared radiation, and the like can also be used. Among the markers that can be selected are auxotrophic markers including, but not limited to, *URA3*, *HIS3*, *TRP1*, *LEU2*, *LYS2*, *ADE2*, and the like. Typically, for purposes of auxotrophic selection, the yeast host cells transformed with bait vector and/or prey vector are cultured in a medium lacking a particular nutrient. Other selectable markers are not based on auxotrophies, but rather on resistance or sensitivity to an antibiotic or other xenobiotic. Examples of such markers include but are not limited to chloramphenicol acetyl transferase (CAT) gene, which

confers resistance to chloramphenicol; *CAN1* gene, which encodes an arginine permease and thereby renders cells sensitive to canavanine (see Sikorski *et al.*, *Meth. Enzymol.*, 194:302-318 (1991)); the bacterial kanamycin resistance gene (kan^R), which renders eukaryotic cells resistant to the aminoglycoside G418 (see Wach *et al.*, *Yeast*, 10:1793-1808 (1994)); and *CYH2* gene, which confers sensitivity to cycloheximide (see Sikorski *et al.*, *Meth. Enzymol.*, 194:302-318 (1991)). In addition, the *CUP1* gene, which encodes metallothionein and thereby confers resistance to copper, is also a suitable selection marker. Each of the above selection markers may be used alone or in combination. One or more selection markers can be included in a particular bait or prey vector. The bait vector and prey vector may have the same or different selection markers. In addition, the selection pressure can be placed on the transformed host cells either before or after mating the haploid yeast cells.

As will be apparent, the selection markers used should complement the host strains in which the bait and/or prey vectors are expressed. In other words, when a gene is used as a selection marker gene, a yeast strain lacking the selection marker gene (or having mutation in the corresponding gene) should be used as host cells. Numerous yeast strains or derivative strains corresponding to various selection markers are known in the art. Many of them have been developed specifically for certain yeast two-hybrid systems. The application and optional modification of such strains with respect to the present invention should be apparent to a skilled artisan apprised of the present disclosure. Methods for genetically manipulating yeast strains using genetic crossing or recombinant mutagenesis are well known in the art. See *e.g.*, Rothstein, *Meth. Enzymol.*, 101:202-211 (1983). By way of example, the following yeast strains are well known in the art, and can be used in the present invention upon necessary modifications and adjustment:

L40 strain which has the genotype *MATa his3 Δ 200 trp1-901 leu2-3,112 ade2 LYS2::(lexAop)4-HIS3 URA3::(lexAop)8-lacZ*;

EGY48 strain which has the genotype *MATa trp1 his3 ura3 6ops-LEU2*; and

MaV103 strain which has the genotype *MATa ura3-52 leu2-3,112 trp1-901 his3 Δ 200 ade2-101 gal4 Δ gal80 Δ SPAL10::URA3 GAL1::HIS3::lys2* (see Kumar *et al.*, *J. Biol. Chem.* 272:13548-13554 (1997); Vidal *et al.*, *Proc. Natl. Acad. Sci. USA*,

93:10315-10320 (1996)). Such strains are generally available in the research community, and can also be obtained by simple yeast genetic manipulation. *See, e.g., The Yeast Two-Hybrid System*, Bartel and Fields, eds., pages 173-182, Oxford University Press, New York, NY, 1997.

5 In addition, the following yeast strains are commercially available:

Y190 strain which is available from Clontech, Palo Alto, CA and has the genotype *MATa gal4 gal80 his3Δ200 trp1-901 ade2-101 ura3-52 leu2-3, 112 URA3::GAL1-lacZ LYS2::GAL1-HIS3 cyh^r*; and

10 YRG-2 Strain which is available from Stratagene, La Jolla, CA and has the genotype *MATa ura3-52 his3-200 ade2-101 lys2-801 trp1-901 leu2-3, 112 gal4-542 gal80-538 LYS2::GAL1-HIS3 URA3::GAL1/CYC1-lacZ*.

15 In fact, different versions of vectors and host strains specially designed for yeast two-hybrid system analysis are available in kits from commercial vendors such as Clontech, Palo Alto, CA and Stratagene, La Jolla, CA, all of which can be modified for use in the present invention.

5.3.1.2. Reporters

20 Generally, in a transcription-based two-hybrid assay, the interaction between a bait fusion protein and a prey fusion protein brings the DNA-binding domain and the transcription-activation domain into proximity forming a functional transcriptional factor that acts on a specific promoter to drive the expression of a reporter protein. The transcription activation domain and the DNA-binding domain may be selected from various known transcriptional activators, e.g., GAL4, GCN4, ARD1, the human estrogen receptor, *E. coli* LexA protein, herpes simplex virus VP16 (Triezenberg *et al.*, *Genes* 25 *Dev.* 2:718-729 (1988)), the *E. coli* B42 protein (acid blob, *see* Gyuris *et al.*, *Cell*, 75:791-803 (1993)), NF-kB p65, and the like. The reporter gene and the promoter driving its transcription typically are incorporated into a separate reporter vector. Alternatively, the host cells are engineered to contain such a promoter-reporter gene sequence in their chromosomes. Thus, the interaction or lack of interaction between two 30 interacting protein members of a protein complex can be determined by detecting or measuring changes in the reporter in the assay system. Although the reporters and

selection markers can be of similar types and used in a similar manner in the present invention, the reporters and selection markers should be carefully selected in a particular detection assay such that they are distinguishable from each other and do not interfere with each other's function.

5 Many different types reporters are useful in the screening assays. For example, a reporter protein may be a fusion protein having an epitope tag fused to a protein.

Commonly used and commercially available epitope tags include sequences derived from, e.g., influenza virus hemagglutinin (HA), Simian Virus 5 (V5), polyhistidine (6xHis), *c-myc*, *lacZ*, GST, and the like. Antibodies specific to these epitope tags are
10 generally commercially available. Thus, the expressed reporter can be detected using an epitope-specific antibody in an immunoassay.

In another embodiment, the reporter is selected such that it can be detected by a color-based assay. Examples of such reporters include, e.g., the *lacZ* protein (β -galactosidase), the green fluorescent protein (GFP), which can be detected by
15 fluorescence assay and sorted by flow-activated cell sorting (FACS) (*See* Cubitt *et al.*, *Trends Biochem. Sci.*, 20:448-455 (1995)), secreted alkaline phosphatase, horseradish peroxidase, the blue fluorescent protein (BFP), and luciferase photoproteins such as aequorin, obelin, mnemiopsin, and berovin (*See* U.S. Patent No. 6,087,476, which is incorporated herein by reference).

20 Alternatively, an auxotrophic factor is used as a reporter in a host strain deficient in the auxotrophic factor. Thus, suitable auxotrophic reporter genes include, but are not limited to, *URA3*, *HIS3*, *TRP1*, *LEU2*, *LYS2*, *ADE2*, and the like. For example, yeast cells containing a mutant *URA3* gene can be used as host cells (Ura⁻ phenotype). Such cells lack *URA3*-encoded functional orotidine-5'-phosphate decarboxylase, an enzyme
25 required by yeast cells for the biosynthesis of uracil. As a result, the cells are unable to grow on a medium lacking uracil. However, wild-type orotidine-5'-phosphate decarboxylase catalyzes the conversion of a non-toxic compound 5-fluoroorotic acid (5-FOA) to a toxic product, 5-fluorouracil. Thus, yeast cells containing a wild-type *URA3* gene are sensitive to 5-FOA and cannot grow on a medium containing 5-FOA.

30 Therefore, when the interaction between the interacting protein members in the fusion proteins results in the expression of active orotidine-5'-phosphate decarboxylase, the Ura⁻

(Foa^R) yeast cells will be able to grow on a uracil deficient medium (SC-Ura plates). However, such cells will not survive on a medium containing 5-FOA. Thus, protein-protein interactions can be detected based on cell growth.

Additionally, antibiotic resistance reporters can also be employed in a similar manner. In this respect, host cells sensitive to a particular antibiotic are used. Antibiotics resistance reporters include, for example, the chloramphenicol acetyl transferase (CAT) gene and the kan^R gene, which confer resistance to G418 in eukaryotes, and kanamycin in prokaryotes, respectively.

5.3.1.3. Screening Assays for Interaction Antagonists

The screening assays of the present invention are useful in identifying compounds capable of interfering with or disrupting or dissociating protein-protein interactions between survivin, or a mutant form thereof, and a protein selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT, or a mutant form thereof. For example, survivin, or a mutant form thereof, and its interacting partners, or mutant forms thereof, are believed to play a role in apoptosis, and thus are involved in cancer, viral infection, AIDS, ischemia, stroke, autoimmune diseases, neurodegenerative diseases, inflammatory disorders, sepsis, and osteoporosis. It may be possible to ameliorate or alleviate the diseases or disorders in a patient by interfering with or dissociating normal interactions between survivin and one of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT. Alternatively, if the disease or disorder is associated with increased expression of survivin and/or one of the survivin-interacting proteins in accordance with the present invention, then the disease may be treated or prevented by weakening or dissociating the interaction between survivin and the survivin-interacting protein in patients. In addition, if a disease or disorder is associated with mutant forms of survivin and/or one of the survivin-interacting proteins that lead to strengthened protein-protein interaction therebetween, then the disease or disorder may be treated with a compound that weakens or interferes with the interaction between the mutant form of survivin and/or the survivin-interacting protein(s).

In a screening assay for an interaction antagonist, survivin (or a homologue, fragment or derivative thereof), or a mutant form of survivin (or a homologue, fragment

or derivative thereof), and a survivin-interacting protein (or a homologue, fragment or derivative thereof), or a mutant form of a survivin-interacting protein (or a homologue, fragment or derivative thereof), are used as test proteins expressed in the form of fusion proteins as described above for purposes of a two-hybrid assay. The fusion proteins are
5 expressed in a host cell and allowed to interact with each other in the presence of one or more test compounds.

In a preferred embodiment, a counterselectable marker is used as a reporter such that a detectable signal (e.g., appearance of color or fluorescence, or cell survival) is present only when the test compound is capable of interfering with the interaction
10 between the two test proteins. In this respect, the reporters used in various “reverse two-hybrid systems” known in the art may be employed. Reverse two-hybrid systems are disclosed in, e.g., U.S. Patent Nos. 5,525,490; 5,733,726; 5,885,779; Vidal *et al.*, *Proc. Natl. Acad. Sci. USA*, 93:10315-10320 (1996); and Vidal *et al.*, *Proc. Natl. Acad. Sci. USA*, 93:10321-10326 (1996), all of which are incorporated herein by reference.

15 Examples of suitable counterselectable reporters useful in a yeast system include the *URA3* gene (encoding orotidine-5'-decarboxylase, which converts 5-fluoroorotic acid (5-FOA) to the toxic metabolite 5-fluorouracil), the *CAN1* gene (encoding arginine permease, which transports toxic arginine analog canavanine into yeast cells), the *GAL1* gene (encoding galactokinase, which catalyzes the conversion of 2-deoxygalactose to
20 toxic 2-deoxygalactose-1-phosphate), the *LYS2* gene (encoding α -aminoadipate reductase, which renders yeast cells unable to grow on a medium containing α -aminoadipate as the sole nitrogen source), the *MET15* gene (encoding O-acetylhomoserine sulphydrylase, which confers on yeast cells sensitivity to methyl mercury), and the *CYH2* gene (encoding L29 ribosomal protein, which confers sensitivity
25 to cycloheximide). In addition, any known cytotoxic agents including cytotoxic proteins such as the diphtheria toxin (DTA) catalytic domain can also be used as counterselectable reporters. See U.S. Patent No. 5,733,726. DTA causes the ADP-ribosylation of elongation factor-2 and thus inhibits protein synthesis and causes cell death. Other examples of cytotoxic agents include ricin, Shiga toxin, and exotoxin A of *Pseudomonas*
30 *aeruginosa*.

For example, when the *URA3* gene is used as a counterselectable reporter gene, yeast cells containing a mutant *URA3* gene can be used as host cells (Ura⁻ Foa^R phenotype) for the *in vivo* assay. Such cells lack *URA3*-encoded functional orotidine-5'-phosphate decarboxylase, an enzyme required for the biosynthesis of uracil. As a result, the cells are unable to grow on media lacking uracil. However, because of the absence of a wild-type orotidine-5'-phosphate decarboxylase, the yeast cells cannot convert non-toxic 5-fluoroorotic acid (5-FOA) to a toxic product, 5-fluorouracil. Thus, such yeast cells are resistant to 5-FOA and can grow on a medium containing 5-FOA. Therefore, for example, to screen for a compound capable of disrupting interactions between survivin (or a homologue, fragment or derivative thereof), or a mutant form of survivin (or a homologue, fragment or derivative thereof), and HDLC1 (or a homologue, fragment or derivative thereof), or a mutant form of HDLC1 (or a homologue, fragment or derivative thereof), survivin (or a homologue, fragment or derivative thereof) can be expressed as a fusion protein with a DNA-binding domain of a suitable transcription activator while HDLC1 (or a homologue, fragment or derivative thereof) is expressed as a fusion protein with a transcription activation domain of a suitable transcription activator. In the host strain, the reporter *URA3* gene may be operably linked to a promoter specifically responsive to the association of the transcription activation domain and the DNA-binding domain. After the fusion proteins are expressed in the Ura⁻ Foa^R yeast cells, an *in vivo* screening assay can be conducted in the presence of a test compound with the yeast cells being cultured on a medium containing uracil and 5-FOA. If the test compound does not disrupt the interaction between survivin and HDLC1, active *URA3* gene product, i.e., orotidine-5'-decarboxylase, which converts 5-FOA to toxic 5-fluorouracil, is expressed. As a result, the yeast cells cannot grow. On the other hand, when the test compound disrupts the interaction between survivin and HDLC1, no active orotidine-5'-decarboxylase is produced in the host yeast cells. Consequently, the yeast cells will survive and grow on the 5-FOA-containing medium. Therefore, compounds capable of interfering with or dissociating the interaction between survivin and HDLC1 can thus be identified based on colony formation.

As will be apparent, the screening assay of the present invention can be applied in a format appropriate for large-scale screening. For example, combinatorial technologies

can be employed to construct combinatorial libraries of small organic molecules or small peptides. *See generally, e.g., Kenan et al., Trends Biochem. Sc., 19:57-64 (1994); Gallop et al., J. Med. Chem., 37:1233-1251 (1994); Gordon et al., J. Med. Chem., 37:1385-1401 (1994); Ecker et al., Biotechnology, 13:351-360 (1995).* Such combinatorial libraries of compounds can be applied to the screening assay of the present invention to isolate specific modulators of particular protein-protein interactions. In the case of random peptide libraries, the random peptides can be co-expressed with the fusion proteins of the present invention in host cells and assayed *in vivo*. *See e.g., Yang et al., Nucl. Acids Res., 23:1152-1156 (1995).* Alternatively, they can be added to the culture medium for uptake by the host cells.

Conveniently, yeast mating is used in an *in vivo* screening assay. For example, haploid cells of a-mating type expressing one fusion protein as described above are mated with haploid cells of α -mating type expressing the other fusion protein. Upon mating, the diploid cells are spread on a suitable medium to form a lawn. Drops of test compounds can be deposited onto different areas of the lawn. After culturing the lawn for an appropriate period of time, drops containing a compound capable of modulating the interaction between the particular test proteins in the fusion proteins can be identified by stimulation or inhibition of growth in the vicinity of the drops.

The screening assays of the present invention for identifying compounds capable of modulating protein-protein interactions can also be fine-tuned by various techniques to adjust the thresholds or sensitivity of the positive and negative selections. Mutations can be introduced into the reporter proteins to adjust their activities. The uptake of test compounds by the host cells can also be adjusted. For example, yeast high uptake mutants such as the *erg6* mutant strains can facilitate yeast uptake of the test compounds. *See Gaber et al., Mol. Cell. Biol., 9:3447-3456 (1989).* Likewise, the uptake of the selection compounds such as 5-FOA, 2-deoxygalactose, cycloheximide, α -aminoadipate, and the like can also be fine-tuned.

5.3.1.4. Screening Assays for Interaction Agonists

The screening assays of the present invention can also be used in identifying compounds that trigger or initiate, enhance or stabilize protein-protein interactions

between survivin, or a mutant form thereof, and a protein selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT, or a mutant thereof. For example, if a disease or disorder is associated with decreased expression of survivin and/or a member of selected from the group of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT, then the disease or disorder may be treated or prevented by strengthening or stabilizing the interaction between survivin and the survivin-interacting member in patients. Alternatively, if a disease or disorder is associated with mutant forms of survivin and/or mutant forms of a survivin-interacting protein that lead to weakened or abolished protein-protein interaction therebetween, then the disease or disorder may be treated with a compound that initiates or stabilizes the interaction between the mutant forms of survivin and/or the mutant forms of survivin-interacting protein(s).

Thus, a screening assay can be performed in the same manner as described above, except that a positively selectable marker is used. For example, survivin (or a homologue, fragment, or derivative thereof), or a mutant form of survivin (or a homologue, fragment, or derivative thereof), and a protein selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT (or a homologue, fragment, or derivative thereof), or a mutant form of a protein selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT (or a homologue, fragment, or derivative thereof), are used as test proteins expressed in the form of fusion proteins as described above for purposes of a two-hybrid assay. The fusion proteins are expressed in host cells and are allowed to interact with each other in the presence of one or more test compounds.

A gene encoding a positively selectable marker such as the lacZ protein may be used as a reporter gene such that when a test compound enables or enhances the interaction between survivin (or a homologue, fragment, or derivative thereof), or a mutant form of survivin (or a homologue, fragment, or derivative thereof), and a protein selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT (or a homologue, fragment, or derivative thereof), or a mutant form of a protein selected from the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT (or a homologue, fragment, or derivative thereof),

the lacZ protein, i.e., β -galactosidase, is expressed. As a result, the compound may be identified based on the appearance of a blue color when the host cells are cultured in a medium containing X-Gal.

Generally, a control assay is performed in which the above screening assay is conducted in the absence of the test compound. The result is then compared with that obtained in the presence of the test compound.

5.4. Optimization of the Identified Compounds

Once the test compounds are selected capable of modulating the interaction between HDLC1 and a protein selected from HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT, or modulating HDLC1, or HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT, a data set including data defining the identity or characteristics of the test compounds can be generated. The data set may include information relating to the properties of a selected test compound, e.g., chemical structure, chirality, molecular weight, melting point, etc. Alternatively, the data set may simply include assigned identification numbers understood by the researchers conducting the screening assay and/or researchers receiving the data set as representing specific test compounds. The data or information can be cast in a transmittable form that can be communicated or transmitted to other researchers, particularly researchers in a different country. Such a transmittable form can vary and can be tangible or intangible. For example, the data set defining one or more selected test compounds can be embodied in texts, tables, diagrams, molecular structures, photographs, charts, images or any other visual forms. The data or information can be recorded on a tangible media such as paper or embodied in computer-readable forms (e.g., electronic, electromagnetic, optical or other signals). The data in a computer-readable form can be stored in a computer usable storage medium (e.g., floppy disks, magnetic tapes, optical disks, and the like) or transmitted directly through a communication infrastructure. In particular, the data embodied in electronic signals can be transmitted in the form of email or posted on a website on the Internet or Intranet. In addition, the information or data on a selected test compound can also be recorded in an audio form and transmitted through any suitable

media, e.g., analog or digital cable lines, fiber optic cables, etc., via telephone, facsimile, wireless mobile phone, Internet phone and the like.

Thus, the information and data on a test compound selected in a screening assay described above or by virtual screening as discussed below can be produced anywhere in the world and transmitted to a different location. For example, when a screening assay is conducted offshore, the information and data on a selected test compound can be generated and cast in a transmittable form as described above. The data and information in a transmittable form thus can be imported into the U.S. or transmitted to any other countries, where the data and information may be used in further testing the selected test compound and/or in modifying and optimizing the selected test compound to develop lead compounds for testing in clinical trials.

Compounds can also be selected based on structural models of the target protein or protein complex and/or test compounds. In addition, once an effective compound is identified, structural analogs or mimetics thereof can be produced based on rational drug design with the aim of improving drug efficacy and stability, and reducing side effects. Methods known in the art for rational drug design can be used in the present invention. *See, e.g., Hodgson et al., Bio/Technology*, 9:19-21 (1991); U.S. Patent Nos. 5,800,998 and 5,891,628, all of which are incorporated herein by reference. An example of rational drug design is the development of HIV protease inhibitors. *See Erickson et al., Science*, 249:527-533 (1990).

In this respect, structural information on the target protein or protein complex is obtained. Preferably, atomic coordinates defining a three-dimensional structure of the target protein or protein complex can be obtained. For example, each of the interacting pair can be expressed and purified. The purified interacting protein pairs are then allowed to interact with each other *in vitro* under appropriate conditions. Optionally, the interacting protein complex can be stabilized by crosslinking or other techniques. The interacting complex can be studied using various biophysical techniques including, e.g., X-ray crystallography, NMR, computer modeling, mass spectrometry, and the like. Likewise, structural information can also be obtained from protein complexes formed by interacting proteins and a compound that initiates or stabilizes the interaction of the proteins. Methods for obtaining such atomic coordinates by X-ray crystallography,

NMR, and the like are known in the art and the application thereof to the target protein or protein complex of the present invention should be apparent to skilled persons in the art of structural biology. See Smyth and Martin, *Mol. Pathol.*, 53:8-14 (2000); Oakley and Wilce, *Clin. Exp. Pharmacol. Physiol.*, 27(3):145-151 (2000); Ferentz and Wagner, *Q. Rev. Biophys.*, 33:29-65 (2000); Hicks, *Curr. Med. Chem.*, 8(6):627-650 (2001); and Roberts, *Curr. Opin. Biotechnol.*, 10:42-47 (1999).

In addition, understanding of the interaction between the proteins of interest in the presence or absence of a modulator can also be derived from mutagenesis analysis using yeast two-hybrid system or other methods for detection protein-protein interaction. In this respect, various mutations can be introduced into the interacting proteins and the effect of the mutations on protein-protein interaction examined by a suitable method such as the yeast two-hybrid system.

Various mutations including amino acid substitutions, deletions and insertions can be introduced into a protein sequence using conventional recombinant DNA technologies. Generally, it is particularly desirable to decipher the protein binding sites. Thus, it is important that the mutations introduced only affect protein-protein interaction and cause minimal structural disturbances. Mutations are preferably designed based on knowledge of the three-dimensional structure of the interacting proteins. Preferably, mutations are introduced to alter charged amino acids or hydrophobic amino acids exposed on the surface of the proteins, since ionic interactions and hydrophobic interactions are often involved in protein-protein interactions. Alternatively, the "alanine scanning mutagenesis" technique is used. See Wells, *et al.*, *Methods Enzymol.*, 202:301-306 (1991); Bass *et al.*, *Proc. Natl. Acad. Sci. USA*, 88:4498-4502 (1991); Bennet *et al.*, *J. Biol. Chem.*, 266:5191-5201 (1991); Diamond *et al.*, *J. Virol.*, 68:863-876 (1994). Using this technique, charged or hydrophobic amino acid residues of the interacting proteins are replaced by alanine, and the effect on the interaction between the proteins is analyzed using e.g., the yeast two-hybrid system. For example, the entire protein sequence can be scanned in a window of five amino acids. When two or more charged or hydrophobic amino acids appear in a window, the charged or hydrophobic amino acids are changed to alanine using standard recombinant DNA techniques. The thus mutated proteins are used as "test proteins" in the above-described two-hybrid assays to examine

the effect of the mutations on protein-protein interaction. Preferably, the mutagenesis analysis is conducted both in the presence and in the absence of an identified modulator compound. In this manner, the domains or residues of the proteins important to protein-protein interaction and/or the interaction between the modulator compound and the

5 interacting proteins can be identified.

Based on the information obtained, structural relationships between the interacting proteins, as well as between the identified modulators and the interacting proteins are elucidated. For example, for the identified modulators (i.e., lead compounds), the three-dimensional structure and chemical moieties critical to their
10 modulating effect on the interacting proteins are revealed. Using this information and various techniques known in the art of molecular modeling (i.e., simulated annealing), medicinal chemists can then design analog compounds that might be more effective modulators of the protein-protein interactions of the present invention. For example, the analog compounds might show more specific or tighter binding to their targets, and
15 thereby might exhibit fewer side effects, or might have more desirable pharmacological characteristics (e.g., greater solubility).

In addition, if the lead compound is a peptide, it can also be analyzed by the alanine scanning technique and/or the two-hybrid assay to determine the domains or residues of the peptide important to its modulating effect on particular protein-protein
20 interactions. The peptide compound can be used as a lead molecule for rational design of small organic molecules or peptide mimetics. See Huber *et al.*, *Curr. Med. Chem.*, 1:13-34 (1994).

The domains, residues or moieties critical to the modulating effect of the identified compound constitute the active region of the compound known as its
25 "pharmacophore." Once the pharmacophore has been elucidated, a structural model can be established by a modeling process that may incorporate data from NMR analysis, X-ray diffraction data, alanine scanning, spectroscopic techniques and the like. Various techniques including computational analysis (e.g., molecular modeling and simulated annealing), similarity mapping and the like can all be used in this modeling process. See
30 e.g., Perry *et al.*, in *OSAR: Quantitative Structure-Activity Relationships in Drug Design*, pp.189-193, Alan R. Liss, Inc., 1989; Rotivinen *et al.*, *Acta Pharmaceutica Fennica*,

97:159-166 (1988); Lewis *et al.*, *Proc. R. Soc. Lond.*, 236:125-140 (1989); McKinaly *et al.*, *Annu. Rev. Pharmacol. Toxicol.*, 29:111-122 (1989). Commercial molecular modeling systems available from Polygen Corporation, Waltham, MA, include the CHARMM program, which performs energy minimization and molecular dynamics
5 functions, and QUANTA program which performs construction, graphic modeling and analysis of molecular structure. Such programs allow interactive construction, modification and visualization of molecules. Other computer modeling programs are also available from BioDesign, Inc. (Pasadena, CA.), Hypercube, Inc. (Cambridge, Ontario), and Allelix, Inc. (Mississauga, Ontario, Canada).

10 A template can be formed based on the established model. Various compounds can then be designed by linking various chemical groups or moieties to the template. Various moieties of the template can also be replaced. In addition, in the case of a peptide lead compound, the peptide or mimetics thereof can be cyclized, e.g., by linking the N-terminus and C-terminus together, to increase its stability. These rationally
15 designed compounds are further tested. In this manner, pharmacologically acceptable and stable compounds with improved efficacy and reduced side effect can be developed. The compounds identified in accordance with the present invention can be incorporated into a pharmaceutical formulation suitable for administration to an individual.

In addition, the structural models or atomic coordinates defining a three-
20 dimensional structure of the target protein or protein complex can also be used in virtual screen to select compounds capable of modulating the target protein or protein complex. Various methods of computer-based virtual screen using atomic coordinates are generally known in the art. For example, U.S. Patent No. 5,798,247 (which is incorporated herein by reference) discloses a method of identifying a compound (specifically, an interleukin
25 converting enzyme inhibitor) by determining binding interactions between an organic compound and binding sites of a binding cavity within the target protein. The binding sites are defined by atomic coordinates.

6. Therapeutic Applications

30 As described above, the interactions between survivin and the survivin-interacting proteins suggest that these proteins and/or the protein complexes formed by them may be

involved in common biological processes and disease pathways. The protein complexes may mediate the functions of survivin and the survivin-interacting proteins in the biological processes or disease pathways. Thus, one may modulate such biological processes or treat diseases by modulating the functions and activities of survivin, a survivin-interacting protein, and/or a protein complex comprising some combination of these proteins. As used herein, modulating survivin, a survivin-interacting protein, or a protein complex comprising some combination of these proteins means altering (enhancing or reducing) the concentrations or activities of the proteins or protein complexes, e.g., increasing the concentrations of survivin, a survivin-interacting protein or a protein complex comprising some combination of these proteins, enhancing or reducing their biological activities, increasing or decreasing their stability, altering their affinity or specificity to certain other biological molecules, etc. For example, a survivin-containing protein complex of the present invention or its members thereof may be involved in apoptosis. Thus, assays such as those described in Section 4 may be used in determining the effect of an aberration in a particular survivin-containing complex or an interacting member thereof on apoptosis. In addition, it is also possible to determine, using the same assay methods, the presence or absence of an association between a survivin-containing complex or an interacting member thereof and a physiological disorder or disease such as cancer, viral infection, AIDS, ischemia, stroke, autoimmune diseases, neurodegenerative diseases, inflammatory disorders, sepsis, and osteoporosis or predisposition to a physiological disorder or disease.

Once such associations are established, the diagnostic methods as described in Section 4 can be used in diagnosing the disease or disorder, or a patient's predisposition to it. In addition, various *in vitro* and *in vivo* assays may be employed to test the therapeutic or prophylactic efficacies of the various therapeutic approaches described in Sections 6.2 and 6.3 that are aimed at modulating the functions and activities of a particular survivin-containing complex of the present invention, or an interacting member thereof. Similar assays can also be used to test whether the therapeutic approaches described in Sections 6.2 and 6.3 result in the modulation of apoptosis. The cell model or transgenic animal model described in Section 7 may be employed in the *in vitro* and *in vivo* assays.

In accordance with this aspect of the present invention, methods are provided for modulating (promoting or inhibiting) a survivin-containing protein complex or interacting member thereof. The human cells can be in *in vitro* cell or tissue cultures. The methods are also applicable to human cells in a patient.

5 In one embodiment, the concentration of a survivin-containing protein complex of the present invention is reduced in the cells. Various methods can be employed to reduce the concentration of the protein complex. The protein complex concentration can be reduced by interfering with the interactions between the interacting members. For example, compounds capable of interfering with interactions between survivin and a
10 protein selected from the group of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT can be administered to the cells *in vitro* or *in vivo* in a patient. Such compounds can be compounds capable of binding survivin or the protein selected from HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT. They can also be antibodies immunoreactive with the survivin or the protein selected from HDLC1,
15 beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT. Also, the compounds can be small peptides derived from the a survivin-interacting protein or mimetics thereof capable of binding survivin, or small peptides derived from survivin protein or mimetics thereof capable of binding a protein selected from HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT.

20 In another embodiment, the method of modulating the protein complex includes inhibiting the expression of survivin protein and/or a survivin-interacting protein. The inhibition can be at the transcriptional, translational, or post-translational level. For example, antisense compounds and ribozyme compounds can be administered to human cells in cultures or in human bodies. In addition, RNA interference technologies may
25 also be employed to administer to cells double-stranded RNA or RNA hairpins capable of "knocking down" the expression of survivin protein and/or a survivin-interacting protein.

In the various embodiments described above, preferably the concentrations or activities of both survivin protein and a survivin-interacting protein are reduced or inhibited.

30 In yet another embodiment, an antibody selectively immunoreactive with a protein complex having survivin interacting with a protein selected from HDLC1, beta-

actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT is administered to cells in vitro or in human bodies to inhibit the protein complex activities and/or reduce the concentration of the protein complex in the cells or patient.

6.1. Applicable Diseases

Apoptosis, also known as programmed cell death, is an active process essential for normal development and functions of multicellular organisms. Typically, apoptosis involves isolated single cells and is characterized by DNA fragmentation, morphological changes of cells and nuclei including cell shrinkage, cell surface blebbing, exposure of phosphatidylserine on the cell surface, involution, contraction, chromatin condensation and fragmentation, and phagocytosis without cell infiltration or inflammation. *See* Honig and Rosenberg, *Am. J. Med.*, 108:317-330 (2000). These characteristics are typically used as markers for assaying apoptosis and can be used in the cell-based assays described in Section 7 below. Many techniques have been developed in the art for detecting such apoptosis markers including, e.g., examining DNA ladders, detecting free DNA ends or breaks under TdT-mediated dUTP nick end labeling (TUNEL) or in situ end labeling (ISEL), determining chromatin clumping by bisbenzimidazole stain or acridine orange stain, observation under light or electron microscopy, immunochemistry analysis of apoptosis-specific proteins, Western blot analysis of caspase-3 cleavage, etc.

Dysregulation of apoptosis can lead to various diseases and disorders. It is now well-known that reduced apoptosis may contribute to tumorigenesis and formation of cancer. Thus, induction of tumor cell apoptosis can be an effective approach in treating cancer. In addition, stimulation of endothelial cell apoptosis may prevent tumor blood supply and cause tumor regression. *See* Dimmeler and Zeiher, *Cir. Res.*, 87:434-439 (2000). Dysregulation of apoptosis is also an integral part of a wide range of autoimmune diseases and disorders. *See* Ravirajan *et al.*, *Int. Rev. Immunol.*, 18:563-589 (1999). In addition, many neurological disorders involve apoptosis. During adulthood, there is little normal neuronal cell death. However, neurological diseases, particularly neurodegenerative diseases are often associated with excessive neural cell death. *See* Honig and Rosenberg, *Am. J. Med.*, 108:317-330 (2000). For example, Parkinson's disease is associated with the loss of substantia nigra pars compacta and sympathetic ganglia, while

Alzheimer's disease is characterized with selective cell loss of entorhinal neurons, and hippocampal neurons, cortical neurons. See Honig and Rosenberg, *Am. J. Med.*, 108:317-330 (2000).

Apoptosis also plays an important role in osteoporotic disorders including, but not limited to, postmenopausal osteoporosis, involutional osteoporosis, and glucocorticoid-induced osteoporosis. See Weinstein, *et al.*, *Am. J. Med.*, 108:153-164 (2000). Generally, under normal conditions, the balance between bone formation, bone resorption, bone cell proliferation and apoptosis maintains nearly constant bone mass. The imbalance of such processes leads to abnormal bone remodeling, and thus osteoporosis and other bone-related diseases. It has been suggested that treatment or prevention of osteoporosis may be achieved by promotion of osteoclast apoptosis and prevention of osteoblast apoptosis. See Weinstein, *et al.*, *Am. J. Med.*, 108:153-164 (2000).

Apoptosis also has physiological significance in animal virus infection. See Kyama *et al.*, *Microbes and Infection*, 2:1111-1117 (2000). Apoptosis of cells infected with viruses may slow the viral multiplication process, although animal viruses typically are able to escape apoptosis of the infected cells. However, it has been suggested that apoptosis of the infected cells triggers the phagocytosis of the dying cells by macrophages. This phagocytosis prevents the leakage of toxic substances that are mediators of dysregulated inflammatory reactions. As a result, dysregulated inflammatory reactions are prevented while specific immune response against the viruses are initiated at the viral infection site. See Kyama *et al.*, *Microbes and Infection*, 2:1111-1117 (2000). On the other hand, in the case of HIV infection, viral infection-induced apoptosis of CD4⁺ T cells contributes to the depletion of CD4⁺ T cells and progression of HIV infection and AIDS, which is associated with immunodeficiency. Thus, inhibition of apoptosis of CD4⁺ T cells may be a strategy in preventing or treating HIV infection and AIDS. See Kirschner *et al.*, *JAIDS J. Acq. Imm. Def. Syn.*, 24:352-362 (2000).

Additionally, apoptosis also plays a role in diseases such as ischemic heart disease, stroke, and sepsis. For example, apoptosis-related neuronal cell death after cerebral ischemia may contribute to stroke. See Johnson *et al.*, *J. Neurotrauma.*, 12:843-52 (1995). Thus, inhibition of apoptosis may be an approach in the development of

therapeutic interventions of ischemic stroke. In addition, the inhibition of endothelial cell apoptosis may improve angiogenesis and vasculogenesis in patients with ischemia, and thus may be an effective method for treating ischemia injuries. *See* Dimmeler and Zeiher, *Cir. Res.*, 87:434-439 (2000).

5 The methods for modulating the functions and activities of a survivin-containing protein complex of the present invention, or an interacting member thereof, may be employed to modulate apoptosis. In addition, the methods may also be used in the treatment or prevention of diseases and disorders such as cancer, viral infection, AIDS, ischemia, stroke, autoimmune diseases, neurodegenerative diseases, inflammatory
10 disorders, sepsis, and osteoporosis.

 Thus, diseases and disorders associated with dysregulation of apoptosis or cellular growth and proliferations may be treated or prevented by modulating the functions and activities of survivin, HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-
15 CAT, or the functions and activities of protein complexes comprising survivin and one or more of the survivin-interacting proteins.

 Accordingly, the present invention provides methods for modulating apoptosis by modulating the functions and activities of one or more of the proteins survivin, HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT, or the protein complexes
20 comprising survivin and one or more of the survivin-interacting proteins. The present invention also provides methods for treating or preventing diseases and disorders associated with apoptosis by modulating apoptosis.

 In one aspect of the present invention, methods are provided to treat cancer by inhibiting the protein complexes of the present invention, or by interfering with or
25 inhibiting the interaction between survivin and one or more of the survivin-interacting proteins, or inhibiting the activities of the survivin-interacting proteins. In a specific embodiment, the methods of treating cancer comprises interfering with the interaction between survivin and HDLC1 or inhibiting HDLC1. Such methods can be applicable to a variety of tumors, i.e., abnormal growth, whether cancerous (malignant) or noncancerous (benign), and whether primary tumors or secondary tumors. Such
30 disorders include but are not limited to lung cancers such as bronchogenic carcinoma (e.g., squamous cell carcinoma, small cell carcinoma, large cell carcinoma, and

adenocarcinoma), alveolar cell carcinoma, bronchial adenoma, chondromatous hamartoma (noncancerous), and sarcoma (cancerous); heart tumors such as myxoma, fibromas and rhabdomyomas; bone tumors such as osteochondromas, condromas, chondroblastomas, chondromyxoid fibromas, osteoid osteomas, giant cell tumors, chondrosarcoma, multiple myeloma, osteosarcoma, fibrosarcomas, malignant fibrous histiocytomas, Ewing's tumor (Ewing's sarcoma), and reticulum cell sarcoma; brain tumors such as gliomas (e.g., glioblastoma multiforme), anaplastic astrocytomas, astrocytomas, and oligodendrogliomas, medulloblastomas, chordoma, Schwannomas, ependymomas, meningiomas, pituitary adenoma, pinealoma, osteomas, and hemangioblastomas, craniopharyngiomas, chordomas, germinomas, teratomas, dermoid cysts, and angiomas; various oral cancers; tumors in digestive system such as leiomyoma, epidermoid carcinoma, adenocarcinoma, leiomyosarcoma, stomach adenocarcinomas, intestinal lipomas, intestinal neurofibromas, intestinal fibromas, polyps in large intestine, familial polyposis such as Gardner's syndrome and Peutz-Jeghers syndrome, colorectal cancers (including colon cancer and rectal cancer); liver cancers such as hepatocellular adenomas, hemangioma, hepatocellular carcinoma, fibrolamellar carcinoma, cholangiocarcinoma, hepatoblastoma, and angiosarcoma; kidney tumors such as kidney adenocarcinoma, renal cell carcinoma, hypernephroma, and transitional cell carcinoma of the renal pelvis; bladder cancers; tumors in blood system including acute lymphocytic (lymphoblastic) leukemia, acute myeloid (myelocytic, myelogenous, myeloblastic, myelomonocytic) leukemia, chronic lymphocytic leukemia (e.g., Sézary syndrome and hairy cell leukemia), chronic myelocytic (myeloid, myelogenous, granulocytic) leukemia, Hodgkin's lymphoma, non-Hodgkin's lymphoma, mycosis fungoides, and myeloproliferative disorders (including myeloproliferative disorders are polycythemia vera, myelofibrosis, thrombocythemia, and chronic myelocytic leukemia); skin cancers such as basal cell carcinoma, squamous cell carcinoma, melanoma, Kaposi's sarcoma, and Paget's disease; head and neck cancers; eye-related cancers such as retinoblastoma and intraocular melanocarcinoma; male reproductive system cancers such as benign prostatic hyperplasia, prostate cancer, and testicular cancers (e.g., seminoma, teratoma, embryonal carcinoma, and choriocarcinoma); breast cancer; female reproductive system cancers such as uterus cancer (endometrial carcinoma), cervical

cancer (cervical carcinoma), ovaries (ovarian carcinoma), vulvar carcinoma, vaginal carcinoma, fallopian tube cancer, and hydatidiform mole; thyroid cancer (including papillary, follicular, anaplastic, or medullary cancer); pheochromocytomas (adrenal gland); noncancerous growths of the parathyroid glands; cancerous or noncancerous
5 growths of the pancreas; etc.

In specific embodiments, the therapeutic and prophylactic methods of the present invention are employed to treat or prevent breast cancers, colon cancers, prostate cancers, lung cancers and skin cancers.

The therapeutic and prophylactic methods of the present invention can also be
10 applied to premalignant conditions to prevent or stop the progression of such conditions towards malignancy, or cause regression of the premalignant conditions. Examples of premalignant conditions include hyperplasia, dysplasia, and metaplasia.

Thus, the term “treating cancer” as used herein, specifically refers to administering therapeutic agents to a patient diagnosed of cancer, i.e., having established
15 cancer in the patient, to inhibit the further growth or spread of the malignant cells in the cancerous tissue, and/or to cause the death of the malignant cells. The term “treating cancer” also encompasses treating a patient having premalignant conditions to stop the progression of, or cause regression of, the premalignant conditions.

The methods of the present invention may also be useful in treating or preventing
20 other diseases and disorders caused by abnormal cell proliferation (hyperproliferation or dysproliferation), e.g., keloid, liver cirrhosis, psoriasis, etc. In addition, the methods may also find applications in promoting wound healing, and other cell and tissue growth-related conditions.

In yet another embodiment, the methods for modulating (preferably inhibiting)
25 the functions and activities of survivin-containing complexes or the interacting protein members thereof may be used in treating or preventing autoimmune diseases and disorders including, but not limited to, rheumatoid arthritis, systemic lupus erythematosus (SLE), Sjogren's syndrome, Canale-Smith syndrome, psoriasis, scleroderma, dermatomyositis, polymyositis, Behcet's syndrome, skin-related autoimmune diseases such
30 as bullus pemphigoid, IgA dermatosis, pemphigus vulgaris, pemphigus foliaceus, dermatitis herpetiformis, contact dermatitis, autoimmune alopecia, erythema nodosa, and

epidermolysis bullous aquisita, drug-induced hemotologic autoimmune disorders, autoimmune thrombocytopenic purpura, autoimmune neutropenia, systemic sclerosis, multiple sclerosis, imflammatory demyelinating, diabetes mellitus, autoimmune polyglandular syndromes, vasculitides, Wegener's granulomatosis, Hashimoto's disease, multinodular goitre, Grave's disease, autoimmune encephalomyelitis (EAE), demyelinating diseases, etc. In a specific embodiment, the treatment is achieved by interfering with the interaction between survivin and one or more of interctors (e.g., HDLC1)

The methods for modulating (preferably inhibiting) apoptosis can also be useful in treating neurodegenerative disorders including, but not limited to, Alzheimer's disease, frontotemporal dementia, Parkinson's disease, Huntington's disease, brain trauma, infarction, hemorrhage, amyotrophic lateral sclerosis/Lou Gehrig's disease (ALS), inherited ataxias such as olivopontocerebellar atrophy (spinocerebellar ataxia type 1), and Machado-Joseph disease (spinocerebellar ataxia type 3).

The present invention further relates to treating ischemia and stroke by modulating (preferably promoting) the functions and activities of one or more of the proteins survivin, HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT, or the protein complexes comprising survivin and one or more of the survivin-interacting proteins, to inhibit apoptosis.

The methods for modulating apoptosis in accordance with the present invention may also be used in treating or preventing osteoporotic disorders such as postmenopausal osteoporosis, involutional osteoporosis, and glucocorticoid-induced osteoporosis.

In another aspect, the methods of the present invention may be useful in treating or preventing diseases or disorders associated with viral infection in animals, particularly humans. Such viral infection can be caused by viruses including, HIV, hepatitis A, hepatitis B, hepatitis C, hepatitis D virus, hepatitis E virus, hepatitis G virus, human foamy virus, human herpes viruses (e.g., human herpes virus 1, human herpes virus 2, human herpes virus 4/Epstein Barr virus, human herpes virus 5, human herpes virus 7), human papilloma virus, human parechovirus 2, human T-cell lymphotropic virus, mumps virus, Measles virus, Rubella virus, Semliki Forest virus, West Nile virus, Colorado tick fever virus, foot-and-mouth disease virus, Marburg virus, polyomavirus, TT virus, Lassa

virus, lymphocytic choriomeningitis virus, vesicular stomatitis virus, influenza viruses, human parainfluenza viruses, respiratory syncytial virus, rotavirus, herpes simplex virus, herpes zoster virus, varicella virus, parvovirus, vaccinia virus, Ebola virus, cytomegalovirus, variola virus, encephalitis viruses, adenovirus, echovirus, rhinoviruses, 5 filoviruses, coxsackievirus, coronavirus, HTLV-I, HTLV-II, Dengue viruses, yellow fever virus, regional hemorrhagic fever viruses, molluscum virus, poliovirus, rabiesvirus, etc. Preferably, the treatment is achieved by interfering with the interaction between survivin and one or more of its interactors (e.g., HDLC1).

10 In one specific embodiment, the methods relate to treating or preventing diseases and disorders caused by lentiviruses or retroviruses, particularly HIV infection and AIDS and/or AIDS-related conditions. The methods comprise interfering with the interaction between survivin and an interacting partner thereof according to the present invention, or by inhibiting a protein complex of the present invention or an interacting member thereof.

As used herein, the term "HIV infection" generally encompasses infection of a 15 host animal, particularly a human host, by the human immunodeficiency virus (HIV) family of retroviruses including, but not limited to, HIV I, HIV II, HIV III (a.k.a. HTLV-III, LAV-1, LAV-2), and the like. "HIV" can be used herein to refer to any strains, forms, subtypes, clades and variations in the HIV family. Thus, treating HIV infection will encompass the treatment of a person who is a carrier of any of the HIV family of 20 retroviruses or a person who is diagnosed of active AIDS, as well as the treatment or prophylaxis of the AIDS-related conditions in such persons. A carrier of HIV may be identified by any methods known in the art. For example, a person can be identified as HIV carrier on the basis that the person is anti-HIV antibody positive, or is HIV-positive, or has symptoms of AIDS. That is, "treating HIV infection" should be understood as 25 treating a patient who is at any one of the several stages of HIV infection progression, which, for example, include acute primary infection syndrome (which can be asymptomatic or associated with an influenza-like illness with fevers, malaise, diarrhea and neurologic symptoms such as headache), asymptomatic infection (which is the long latent period with a gradual decline in the number of circulating CD⁴⁺ T cells), and AIDS 30 (which is defined by more serious AIDS-defining illnesses and/or a decline in the circulating CD4 cell count to below a level that is compatible with effective immune

function). In addition, "treating or preventing HIV infection" will also encompass treating suspected infection by HIV after suspected past exposure to HIV by e.g., contact with HIV-contaminated blood, blood transfusion, exchange of body fluids, "unsafe" sex with an infected person, accidental needle stick, receiving a tattoo or acupuncture with contaminated instruments, or transmission of the virus from a mother to a baby during pregnancy, delivery or shortly thereafter. The term "treating HIV infection" may also encompass treating a person who is free of HIV infection but is believed to be at risk of infection by HIV.

The term "treating AIDS" means treating a patient who exhibits more serious AIDS-defining illnesses and/or a decline in the circulating CD4 cell count to below a level that is compatible with effective immune function. The term "treating AIDS" also encompasses treating AIDS-related conditions, which means disorders and diseases incidental to or associated with AIDS or HIV infection such as AIDS-related complex (ARC), progressive generalized lymphadenopathy (PGL), anti-HIV antibody positive conditions, and HIV-positive conditions, AIDS-related neurological conditions (such as dementia or tropical paraparesis), Kaposi's sarcoma, thrombocytopenia purpura and associated opportunistic infections such as *Pneumocystis carinii* pneumonia, *Mycobacterial tuberculosis*, esophageal candidiasis, toxoplasmosis of the brain, CMV retinitis, HIV-related encephalopathy, HIV-related wasting syndrome, etc.

Thus, the term "preventing AIDS" as used herein means preventing in a patient who has HIV infection or is suspected to have HIV infection or is at risk of HIV infection from developing AIDS (which is characterized by more serious AIDS-defining illnesses and/or a decline in the circulating CD4 cell count to below a level that is compatible with effective immune function) and/or AIDS-related conditions.

In another specific embodiment, the present invention provides methods for treating or preventing HBV infection and hepatitis B by interfering with the interaction between survivin and an interacting partner thereof according to the present invention, or by inhibiting a protein complex of the present invention or an interacting member thereof. As used herein, the term "HBV infection" generally encompasses infection of a human by any strain or serotype of hepatitis B virus, including acute hepatitis B infection and chronic hepatitis B infection. Thus, treating HBV infection means the treatment of a

person who is a carrier of any strain or serotype of hepatitis B virus or a person who is diagnosed of active hepatitis B to reduce the HBV viral load in the person or to alleviate one or more symptoms associated with HBV infection and/or hepatitis B, including, e.g., nausea and vomiting, loss of appetite, fatigue, muscle and joint aches, elevated

5 transaminase blood levels, increased prothrombin time, jaundice (yellow discoloration of the eyes and body) and dark urine. A carrier of HBV may be identified by any methods known in the art. For example, a person can be identified as HBV carrier on the basis that the person is anti-HBV antibody positive (e.g., based on hepatitis B core antibody or hepatitis B surface antibody), or is HBV-positive (e.g., based on hepatitis B surface
10 antigens (HBeAg or HbsAg) or HBV RNA or DNA) or has symptoms of hepatitis B infection or hepatitis B. That is, "treating HBV infection" should be understood as treating a patient who is at any one of the several stages of HBV infection progression. In addition, the term "treating HBV infection" will also encompass treating suspected infection by HBV after suspected past exposure to HBV by, e.g., contact with HBV-
15 contaminated blood, blood transfusion, exchange of body fluids, "unsafe" sex with an infected person, accidental needle stick, receiving a tattoo or acupuncture with contaminated instruments, or transmission of the virus from a mother to a baby during pregnancy, delivery or shortly thereafter. The term "treating HBV infection" will also encompass treating a person who is free of HBV infection but is believed to be at risk of
20 infection by HBV.

The term "preventing hepatitis B" as used herein means preventing in a patient who has HBV infection or is suspected to have HBV infection or is at risk of HBV infection from developing hepatitis B (which are characterized by more serious hepatitis-defining symptoms), cirrhosis, or hepatocellular carcinoma.

25 In another specific embodiment, the present invention provides methods for treating or preventing HCV infection and hepatitis C by by interfering with the interaction between survivin and an interacting partner thereof according to the present invention, or by inhibiting a protein complex of the present invention or an interacting member thereof.

30 As used herein, the term "HCV infection" generally encompasses infection of a human by any types or subtypes of hepatitis C virus, including acute hepatitis C infection

and chronic hepatitis C infection. Thus, treating HCV infection means the treatment of a person who is a carrier of any types or subtypes of hepatitis C virus or a person who is diagnosed of active hepatitis C to reduce the HCV viral load in the person or to alleviate one or more symptoms associated with HCV infection and/or hepatitis C. A carrier of HCV may be identified by any methods known in the art. For example, a person can be identified as HCV carrier on the basis that the person is anti-HCV antibody positive, or is HCV-positive (e.g., based on HCV RNA or DNA) or has symptoms of hepatitis C infection or hepatitis C (e.g., elevated serum transaminases). That is, "treating HCV infection" should be understood as treating a patient who is at any one of the several stages of HCV infection progression. In addition, the term "treating HCV infection" will also encompass treating suspected infection by HCV after suspected past exposure to HCV by, e.g., contact with HCV-contaminated blood, blood transfusion, exchange of body fluids, "unsafe" sex with an infected person, accidental needle stick, receiving a tattoo or acupuncture with contaminated instruments, or transmission of the virus from a mother to a baby during pregnancy, delivery or shortly thereafter. The term "treating HCV infection" will also encompass treating a person who is free of HCV infection but is believed to be at risk of infection by HCV. The term of "preventing HCV" as used herein means preventing in a patient who has HCV infection or is suspected to have HCV infection or is at risk of HCV infection from developing hepatitis C (which is characterized by more serious hepatitis-defining symptoms), cirrhosis, or hepatocellular carcinoma.

6.2. Inhibiting Protein Complex or Interacting Protein Members Thereof

In one aspect of the present invention, methods are provided for reducing in cells or tissue the concentration and/or activity of a protein complex identified in accordance with the present invention that comprises survivin and one or more members of the group HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT. In addition, methods are also provided for reducing in cells or tissue the concentration and/or activity of a survivin-interacting protein selected from the group HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT. By reducing the concentration of protein complex and/or the survivin-interacting protein concentration(s) and/or inhibiting the

functional activities of the protein complex and/or the survivin-interacting protein(s), the diseases involving such protein complex or survivin-interacting protein(s) may be treated or prevented.

5

6.2.1. Antibody Therapy

In one embodiment, an antibody may be administered to cells or tissue *in vitro* or to patients. The antibody administered may be immunoreactive with survivin or a member of the group HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT, or protein complexes comprising survivin and a member, or members, of the group HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT. Suitable antibodies may be monoclonal or polyclonal that fall within any antibody class, e.g., IgG, IgM, IgA, etc. The antibody suitable for this invention may also take a form of various antibody fragments including, but not limited to, Fab and F(ab')₂, single-chain fragments (scFv), and the like. In another embodiment, an antibody selectively immunoreactive with the protein complex formed from survivin and one or more survivin-interacting protein, or proteins, in accordance with the present invention is administered to cells or tissue *in vitro* or in a patient. In yet another embodiment, an antibody specific to a survivin-interacting protein selected from the group HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT is administered to cells or tissue *in vitro* or in a patient. Methods for making the antibodies of the present invention should be apparent to a person of skill in the art, especially in view of the discussions in Section 3 above. The antibodies can be administered in any suitable form and route as described in Section 8 below. Preferably, the antibodies are administered in a pharmaceutical composition together with a pharmaceutically acceptable carrier.

25

Alternatively, the antibodies may be delivered by a gene-therapy approach. That is, nucleic acids encoding the antibodies, particularly single-chain fragments (scFv), may be introduced into cells or tissue *in vitro* or in a patient such that desirable antibodies may be produced recombinantly *in vivo* from the nucleic acids. For this purpose, the nucleic acids with appropriate transcriptional and translation regulatory sequences can be directly administered into the patient. Alternatively, the nucleic acids can be incorporated into a suitable vector as described in Sections 2.2 and 5.3.1.1 and delivered into cells or tissue

30

in vitro or in a patient along with the vector. The expression vector containing the nucleic acids can be administered directly to cells or tissue *in vitro* or in a patient. It can also be introduced into cells, preferably cells derived from a patient to be treated, and subsequently delivered into the patient by cell transplantation. See Section 6.3.2 below.

5

6.2.2. Antisense Therapy

In another embodiment, antisense compounds specific to nucleic acids encoding one or more interacting protein members of a protein complex identified in the present invention are administered to cells or tissue *in vitro* or in a patient to be therapeutically or prophylactically treated. The antisense compounds should specifically inhibit the expression of the one or more interacting protein members. As is known in the art, antisense drugs generally act by hybridizing to a particular target nucleic acid thus blocking gene expression. Methods for designing antisense compounds and using such compounds in treating diseases are well known and well developed in the art. For example, the antisense drug Vitravene® (fomivirsen), a 21-base long oligonucleotide, has been successfully developed and marketed by Isis Pharmaceuticals, Inc. for treating cytomegalovirus (CMV)-induced retinitis.

10

15

20

25

30

Any methods for designing and making antisense compounds may be used for purpose of the present invention. See generally, Sanghvi *et al.*, eds., *Antisense Research and Applications*, CRC Press, Boca Raton, 1993. Typically, antisense compounds are oligonucleotides designed based on the nucleotide sequence of the mRNA or gene of one or more target proteins, e.g., the interacting protein members of a particular protein complex of the present invention. In particular, antisense compounds can be designed to specifically hybridize to a particular region of the gene sequence or mRNA of one or more of the interacting protein members to modulate (increase or decrease), replication, transcription, or translation. As used herein, the term “specifically hybridize” or paraphrases thereof means a sufficient degree of complementarity or pairing between an antisense oligo and a target DNA or mRNA such that stable and specific binding occurs therebetween. In particular, 100% complementary or pairing is not required. Specific hybridization takes place when sufficient hybridization occurs between the antisense compound and its intended target nucleic acids in the substantial absence of non-specific

binding of the antisense compound to non-target sequences under predetermined conditions, e.g., for purposes of *in vivo* treatment, preferably under physiological conditions. Preferably, specific hybridization results in the interference with normal expression of the target DNA or mRNA.

5 For example, antisense oligonucleotides can be designed to specifically hybridize to target genes, in regions critical for regulation of transcription; to pre-mRNAs, in regions critical for correct splicing of nascent transcripts; and to mature mRNAs, in regions critical for translation initiation or mRNA stability and localization.

10 As is generally known in the art, commonly used oligonucleotides are oligomers or polymers of ribonucleotides or deoxyribonucleotides, that are composed of a naturally-occurring nitrogenous base, a sugar (ribose or deoxyribose) and a phosphate group. In nature, the nucleotides are linked together by phosphodiester bonds between the 3' and 5' positions of neighboring sugar moieties. However, it is noted that the term "oligonucleotides" also encompasses various non-naturally occurring mimetics and
15 derivatives, i.e., modified forms, of naturally occurring oligonucleotides as described below. Typically an antisense compound of the present invention is an oligonucleotide having from about 6 to about 200, and preferably from about 8 to about 30 nucleoside bases.

20 The antisense compounds preferably contain modified backbones or non-natural internucleoside linkages, including but not limited to, modified phosphorous-containing backbones and non-phosphorous backbones such as morpholino backbones; siloxane, sulfide, sulfoxide, sulfone, sulfonate, sulfonamide, and sulfamate backbones; formacetyl and thioformacetyl backbones; alkene-containing backbones; methyleneimino and methylenehydrazino backbones; amide backbones, and the like.

25 Examples of modified phosphorous-containing backbones include, but are not limited to phosphorothioates, phosphorodithioates, chiral phosphorothioates, phosphotriesters, aminoalkylphosphotriesters, alkyl phosphonates, thionoalkylphosphonates, phosphinates, phosphoramidates, thionophosphoramidates, thionoalkylphosphotriesters, and boranophosphates and various salt forms thereof. *See*
30 e.g., U.S. Pat. Nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496;

5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; and 5,625,050, each of which is herein incorporated by reference.

Examples of the non-phosphorous containing backbones described above are disclosed in, e.g., U.S. Pat. Nos. 5,034,506; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,677,437; and 5,677,439, each of which is herein incorporated by reference.

Another useful modified oligonucleotide is peptide nucleic acid (PNA), in which the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, e.g., an aminoethylglycine backbone. See U.S. Patent Nos. 5,539,082 and 5,714,331; and Nielsen *et al.*, *Science*, 254, 1497-1500 (1991), all of which are incorporated herein by reference. PNA antisense compounds are resistant to RNase H digestion and thus exhibit longer half-life. In addition, various modifications may be made in PNA backbones to impart desirable drug profiles such as better stability, increased drug uptake, higher affinity to target nucleic acid, etc.

Alternatively, the antisense compounds are oligonucleotides containing modified nucleosides, i.e., modified purine or pyrimidine bases, e.g., 5-substituted pyrimidines, 6-azapyrimidines, and N-2, N-6 and O-substituted purines, and the like. See e.g., U.S. Pat. Nos. 3,687,808; 4,845,205; 5,130,302; 5,175,273; 5,367,066; 5,432,272; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,587,469; 5,594,121; 5,596,091; 5,681,941; and 5,750,692, each of which is incorporated herein by reference in its entirety.

In addition, oligonucleotides with substituted or modified sugar moieties may also be used. For example, an antisense compound may have one or more 2'-O-methoxyethyl sugar moieties. See e.g., U.S. Pat. Nos. 4,981,957; 5,118,800; 5,319,080; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,567,811; 5,576,427; 5,591,722; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920, each of which is herein incorporated by reference.

Other types of oligonucleotide modifications are also useful including linking an oligonucleotide to a lipid, phospholipid or cholesterol moiety, cholic acid, thioether, aliphatic chain, polyamine, polyethylene glycol (PEG), or a protein or peptide. The modified oligonucleotides may exhibit increased uptake into cells, and improved

stability, i.e., resistance to nuclease digestion and other biodegradations. *See e.g.*, U.S. Patent No. 4,522,811; Burnham, *Am. J. Hosp. Pharm.*, 15:210-218 (1994).

Antisense compounds can be synthesized using any suitable methods known in the art. In fact, antisense compounds may be custom made by commercial suppliers.

5 Alternatively, antisense compounds may be prepared using DNA synthesizers available commercially from various vendors, e.g., Applied Biosystems Group of Norwalk, CT.

The antisense compounds can be formulated into a pharmaceutical composition with suitable carriers and administered into cells or tissue *in vitro* or in a patient using any suitable route of administration. Alternatively, the antisense compounds may also be
10 used in a “gene-therapy” approach. That is, the oligonucleotide is subcloned into a suitable vector and transformed into human cells. The antisense oligonucleotide is then produced *in vivo* through transcription. Methods for gene therapy are disclosed in Section 6.3.2 below.

15 6.2.3. Ribozyme Therapy

In another embodiment, an enzymatic RNA or ribozyme is designed to target the nucleic acids encoding one or more of the interacting protein members of the protein complex of the present invention. Ribozymes are RNA molecules possessing enzymatic activity. One class of ribozymes is capable of repeatedly cleaving other separate RNA
20 molecules into two or more pieces in a nucleotide base sequence specific manner. *See Kim et al., Proc. Natl. Acad. of Sci. USA*, 84:8788 (1987); Haseloff and Gerlach, *Nature*, 334:585 (1988); and Jefferies *et al., Nucleic Acid Res.*, 17:1371 (1989). Such ribozymes typically have two functional domains: a catalytic domain and a binding sequence that guides the binding of ribozymes to a target RNA through complementary base-pairing.

25 Once a specifically-designed ribozyme is bound to a target mRNA, it enzymatically cleaves the target mRNA, typically reducing its stability and destroying its ability to direct translation of an encoded protein. After a ribozyme has cleaved its RNA target, it is released from that target RNA and thereafter can bind and cleave another target. That is, a single ribozyme molecule can repeatedly bind and cleave new targets. Therefore,
30 one advantage of ribozyme treatment is that a lower amount of exogenous RNA is required as compared to conventional antisense therapies. In addition, ribozymes exhibit

less affinity to mRNA targets than DNA-based antisense oligonucleotides, and therefore are less prone to bind to wrong targets.

In accordance with the present invention, a ribozyme may target any portion of the mRNA of one or more interacting protein members including survivin, and HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT. Methods for selecting a ribozyme target sequence and designing and making ribozymes are generally known in the art. *See e.g.*, U.S. Patent Nos. 4,987,071; 5,496,698; 5,525,468; 5,631,359; 5,646,020; 5,672,511; and 6,140,491, each of which is incorporated herein by reference in its entirety. For example, suitable ribozymes may be designed in various configurations such as hammerhead motifs, hairpin motifs, hepatitis delta virus motifs, group I intron motifs, or RNase P RNA motifs. *See e.g.*, U.S. Patent Nos. 4,987,071; 5,496,698; 5,525,468; 5,631,359; 5,646,020; 5,672,511; and 6,140,491; Rossi *et al.*, *AIDS Res. Human Retroviruses* 8:183 (1992); Hampel and Tritz, *Biochemistry* 28:4929 (1989); Hampel *et al.*, *Nucleic Acids Res.*, 18:299 (1990); Perrotta and Been, *Biochemistry* 31:16 (1992); and Guerrier-Takada *et al.*, *Cell*, 35:849 (1983).

Ribozymes can be synthesized by the same methods used for normal RNA synthesis. For example, such methods are disclosed in Usman *et al.*, *J. Am. Chem. Soc.*, 109:7845-7854 (1987) and Scaringe *et al.*, *Nucleic Acids Res.*, 18:5433-5441 (1990). Modified ribozymes may be synthesized by the methods disclosed in, *e.g.*, U.S. Pat. No. 5,652,094; International Publication Nos. WO 91/03162; WO 92/07065 and WO 93/15187; European Patent Application No. 92110298.4; Perrault *et al.*, *Nature*, 344:565 (1990); Pieken *et al.*, *Science*, 253:314 (1991); and Usman and Cedergren, *Trends in Biochem. Sci.*, 17:334 (1992).

Ribozymes of the present invention may be administered to cells by any known methods, *e.g.*, disclosed in International Publication No. WO 94/02595. For example, they can be administered directly to cells or tissue *in vitro* or in a patient through any suitable route, *e.g.*, intravenous injection. Alternatively, they may be delivered encapsulated in liposomes, by iontophoresis, or by incorporation into other vehicles such as hydrogels, cyclodextrins, biodegradable nanocapsules, and bioadhesive microspheres. In addition, they may also be delivered by gene therapy approach, using a DNA vector

from which the ribozyme RNA can be transcribed directly. Gene therapy methods are disclosed in detail below in Section 6.3.2.

6.2.4. Other Methods

5 The in-patient concentrations and activities of the protein complexes and interacting proteins of the present invention may also be altered by other methods. For example, compounds identified in accordance with the methods described in Section 5 that are capable of interfering with or dissociating protein-protein interactions between the interacting protein members of a protein complex may be administered to cells or
10 tissue *in vitro* or in a patient. Compounds identified in *in vitro* binding assays described in Section 5.2 that bind to the survivin-containing protein complex or the interacting members thereof may also be used in the treatment. Compounds identified in *in vitro* binding assays described in Section 5.2 that bind to the survivin-containing protein complex, or the interacting members thereof, may also be used in the treatment.

15 In addition, potentially useful agents also include incomplete proteins, i.e., fragments of the interacting protein members that are capable of binding to their respective binding partners in a protein complex but are defective with respect to their normal cellular functions. For example, binding domains of the interacting member proteins of a protein complex may be used as competitive inhibitors of the activities of
20 the protein complex. As will be apparent to skilled artisans, derivatives or homologues of the binding domains may also be used. Binding domains can be easily identified using molecular biology techniques, e.g., mutagenesis in combination with yeast two-hybrid assays. Preferably, the protein fragment used is a fragment of an interacting protein member having a length of less than 90%, 80%, more preferably less than 75%, 65%,
25 50%, or less than 40% of the full length of the protein member. In one embodiment, a survivin protein fragment is administered. In a specific embodiment, one or more of the interaction domains of survivin within the regions listed in Table 1 are administered to cells or tissue *in vitro*, or are administered to a patient in need of such treatment. For example, suitable protein fragments can include polypeptides having a contiguous span of
30 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 18, 20 or 25, preferably from 4 to 30, 40 or 50 amino acids or more of the sequence of survivin that are capable of interacting with one

or more proteins selected from the group of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT. Also, suitable protein fragments can also include peptides capable of binding one or more proteins selected from the group of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT and having an amino acid sequence
5 of from 4 to 30 amino acids that is at least 75%, 80%, 82%, 85%, 87%, 90%, 95% or more identical to a contiguous span of amino acids of survivin of the same length.

Alternatively, a polypeptide capable of interacting with survivin and having a contiguous span of 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 18, 20 or 25, preferably from 4 to 30, 40 or 50 or more amino acids of the amino acid sequence of a protein selected from the group
10 of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT may be administered. Also, other examples of suitable compounds include a peptide capable of binding survivin and having an amino acid sequence of from 4 to 30, 40, 50 or more amino acids that is at least 75%, 80%, 82%, 85%, 87%, 90%, 92%, 95% or more identical to a contiguous span of amino acids of the same length from a protein selected
15 from the group of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT. In addition, the administered compounds can be an antibody or antibody fragment, preferably single-chain antibody immunoreactive with survivin or a protein selected from the group of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT, or a protein complex of the present invention.

20 The protein fragments suitable as competitive inhibitors can be delivered into cells by direct cell internalization, receptor mediated endocytosis, or via a "transporter." It is noted that when the target proteins or protein complexes to be modulated reside inside cells, the compound administered to cells *in vitro* or *in vivo* in the method of the present invention preferably is delivered into the cells in order to achieve optimal results.
25 Thus, preferably, the compound to be delivered is associated with a transporter capable of increasing the uptake of the compound by cells harboring the target protein or protein complex. As used herein, the term "transporter" refers to an entity (e.g., a compound or a composition or a physical structure formed from multiple copies of a compound or multiple different compounds) that is capable of facilitating the uptake of a compound of
30 the present invention by animal cells, particularly human cells. Typically, the cell uptake of a compound of the present invention in the presence of a "transporter" is at least 20%

higher, preferably at least 40%, 50%, 75%, and more preferably at least 100% higher than the cell uptake of the compound in the absence of the "transporter."

Many molecules and structures known in the art can be used as "transporters." In one embodiment, a penetratin is used as a transporter. For example, the homeodomain of Antennapedia, a *Drosophila* transcription factor, can be used as a transporter to deliver a compound of the present invention. Indeed, any suitable member of the penetratin class of peptides can be used to carry a compound of the present invention into cells.

Penetratins are disclosed in, e.g., Derossi *et al.*, *Trends Cell Biol.*, 8:84-87 (1998), which is incorporated herein by reference. Penetratins transport molecules attached thereto

across cytoplasmic membranes or nuclear membranes efficiently, in a receptor-independent, energy-independent, and cell type-independent manner. Methods for using a penetratin as a carrier to deliver oligonucleotides and polypeptides are also disclosed in U.S. Patent No. 6,080,724; Pooga *et al.*, *Nat. Biotech.*, 16:857 (1998); and Schutze *et al.*, *J. Immunol.*, 157:650 (1996), all of which are incorporated herein by reference. U.S.

Patent No. 6,080,724 defines the minimal requirements for a penetratin peptide as a peptide of 16 amino acids with 6 to 10 of which being hydrophobic. The amino acid at position 6 counting from either the N- or C-terminus is tryptophan, while the amino acids at positions 3 and 5 counting from either the N- or C-terminus are not both valine.

Preferably, the helix 3 of the homeodomain of *Drosophila* Antennapedia is used as a transporter. More preferably, a peptide having a sequence of amino acid residues 43-58 of the homeodomain Antp is employed as a transporter. In addition, other naturally occurring homologs of the helix 3 of the homeodomain of *Drosophila* Antennapedia can be used. For example, homeodomains of Fushi-tarazu and Engrailed have been shown to be capable of transporting peptides into cells. See Han *et al.*, *Mol. Cells*, 10:728-32

(2000). As used herein, the term "penetratin" also encompasses peptoid analogs of the penetratin peptides. Typically, the penetratin peptides and peptoid analogs thereof are covalently linked to a compound to be delivered into cells thus increasing the cellular uptake of the compound.

In another embodiment, the HIV-1 tat protein or a derivative thereof is used as a "transporter" covalently linked to a compound according to the present invention. The use of HIV-1 tat protein and derivatives thereof to deliver macromolecules into cells has

been known in the art. *See* Green and Loewenstein, *Cell*, 55:1179 (1988); Frankel and Pabo, *Cell*, 55:1189 (1988); Vives *et al.*, *J. Biol. Chem.*, 272:16010-16017 (1997); Schwarze *et al.*, *Science*, 285:1569-1572 (1999). It is known that the sequence responsible for cellular uptake consists of the highly basic region, amino acid residues 49-57. *See e.g.*, Vives *et al.*, *J. Biol. Chem.*, 272:16010-16017 (1997); Wender *et al.*, *Proc. Nat'l Acad. Sci. USA*, 97:13003-13008 (2000). The basic domain is believed to target the lipid bilayer component of cell membranes. It causes a covalently linked protein or nucleic acid to cross cell membrane rapidly in a cell type-independent manner. Proteins ranging in size from 15 to 120 kD have been delivered with this technology into a variety of cell types both *in vitro* and *in vivo*. *See* Schwarze *et al.*, *Science*, 285:1569-1572 (1999). Any HIV tat-derived peptides or peptoid analogs thereof capable of transporting macromolecules such as peptides can be used for purposes of the present invention. For example, any native tat peptides having the highly basic region, amino acid residues 49-57 can be used as a transporter by covalently linking it to the compound to be delivered. In addition, various analogs of the tat peptide of amino acid residues 49-57 can also be useful transporters for purposes of this invention. Examples of various such analogs are disclosed in Wender *et al.*, *Proc. Nat'l Acad. Sci. USA*, 97:13003-13008 (2000) (which is incorporated herein by reference) including, e.g., *d*-Tat₄₉₋₅₇, retro-inverso isomers of *l*- or *d*-Tat₄₉₋₅₇ (i.e., *l*-Tat₅₇₋₄₉ and *d*-Tat₅₇₋₄₉), L-arginine oligomers, D-arginine oligomers, L-lysine oligomers, D-lysine oligomers, L-histidine oligomers, D-histidine oligomers, L-ornithine oligomers, D-ornithine oligomers, and various homologues, derivatives (e.g., modified forms with conjugates linked to the small peptides) and peptoid analogs thereof.

Other useful transporters known in the art include, but are not limited to, short peptide sequences derived from fibroblast growth factor (*See* Lin *et al.*, *J. Biol. Chem.*, 270:14255-14258 (1998)), Galparan (*See* Pooga *et al.*, *FASEB J.* 12:67-77 (1998)), and HSV-1 structural protein VP22 (*See* Elliott and O'Hare, *Cell*, 88:223-233 (1997)).

As the above-described various transporters are generally peptides, fusion proteins can be conveniently made by recombinant expression to contain a transporter peptide covalently linked by a peptide bond to a competitive protein fragment.

Alternatively, conventional methods can be used to chemically synthesize a transporter peptide or a peptide of the present invention or both.

The hybrid peptide can be administered to cells or tissue *in vitro* or to a patient in a suitable pharmaceutical composition as provided in Section 8.

5 In addition to peptide-based transporters, various other types of transporters can also be used, including but not limited to cationic liposomes (*see Rui et al., J. Am. Chem. Soc.*, 120:11213-11218 (1998)), dendrimers (Kono *et al., Bioconjugate Chem.*, 10:1115-1121 (1999)), siderophores (Ghosh *et al., Chem. Biol.*, 3:1011-1019 (1996)), etc. In a specific embodiment, the compound according to the present invention is encapsulated
10 into liposomes for delivery into cells.

 Additionally, when a compound according to the present invention is a peptide, it can be administered to cells by a gene therapy method. That is, a nucleic acid encoding the peptide can be administered to *in vitro* cells or to cells *in vivo* in a human or animal body. Any suitable gene therapy methods may be used for purposes of the present
15 invention. Various gene therapy methods are well known in the art and are described in Section 6.3.2. below. Successes in gene therapy have been reported recently. *See e.g., Kay et al., Nature Genet.*, 24:257-61 (2000); Cavazzana-Calvo *et al., Science*, 288:669 (2000); and Blaese *et al., Science*, 270: 475 (1995); Kantoff, *et al., J. Exp. Med.*, 166:219 (1987).

20 In yet another embodiment, the gene therapy methods discussed in Section 6.3.2 below are used to “knock out” the gene encoding an interacting protein member of a protein complex, or to reduce the gene expression level. For example, the gene may be replaced with a different gene sequence or a non-functional sequence or simply deleted by homologous recombination. In another gene therapy embodiment, the method
25 disclosed in U.S. Patent No. 5,641,670, which is incorporated herein by reference, may be used to reduce the expression of the genes for the interacting protein members. Essentially, an exogenous DNA having at least a regulatory sequence, an exon and a splice donor site can be introduced into an endogenous gene encoding an interacting protein member by homologous recombination such that the regulatory sequence, the
30 exon and the splice donor site present in the DNA construct become operatively linked to the endogenous gene. As a result, the expression of the endogenous gene is controlled by

the newly introduced exogenous regulatory sequence. Therefore, when the exogenous regulatory sequence is a strong gene expression repressor, the expression of the endogenous gene encoding the interacting protein member is reduced or blocked. *See* U.S. Patent No. 5,641,670.

5

6.3. Activation of Protein Complex or Interacting Protein Members Thereof

The present invention also provides methods for increasing in cells or tissue *in vitro* or in a patient the concentration and/or activity of a protein complex, or of an individual protein member thereof, identified in accordance with the present invention.

10 Such methods can be particularly useful in instances where a reduced concentration and/or activity of a protein complex, or a protein member thereof, is associated with a particular disease or disorder to be treated, or where an increased concentration and/or activity of a protein complex, or a protein member thereof, would be beneficial to the improvement of a cellular function or disease state. By increasing the concentration of
15 the protein complex, or a protein member thereof, and/or stimulating the functional activities of the protein complex or a protein member thereof, the disease or disorder may be treated or prevented.

6.3.1. Administration of Protein Complex or Protein Members Thereof

20 Where the concentration or activity of a particular survivin-containing protein complex, or survivin itself, or a survivin-interacting protein of the present invention, in cells or tissue *in vitro* or in a patient is determined to be low or is desired to be increased, the protein complex, or survivin, or the survivin-interacting protein may be administered directly to the patient to increase the concentration and/or activity of the protein complex,
25 survivin, or the survivin-interacting protein. For this purpose, protein complexes prepared by any one of the methods described in Section 2.2 may be administered to the patient, preferably in a pharmaceutical composition as described below. Alternatively, one or more individual interacting protein members of the protein complex may also be administered to the patient in need of treatment. For example, one or more proteins such
30 as survivin, HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT may be given to cells or tissue *in vitro* or to a patient. Proteins isolated or purified from

normal individuals or recombinantly produced can all be used in this respect. Preferably, two or more interacting protein members of a protein complex are administered. The proteins or protein complexes may be administered to a patient needing treatment using any of the methods described in Section 8.

5

6.3.2. Gene Therapy

In another embodiment, the concentration and/or activity of a particular survivin-containing protein complex or survivin, or a known survivin-interacting protein (selected from the group including HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT) is increased or restored in patients, tissue or cells by a gene therapy approach. For example, nucleic acids encoding one or more protein members of a survivin-containing protein complex of the present invention, or portions or fragments thereof are introduced into patients, tissue, or cells such that the protein(s) are expressed from the introduced nucleic acids. For these purposes, nucleic acids encoding one or more of survivin, HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT, or fragments, homologues or derivatives thereof can be used in the gene therapy in accordance with the present invention. For example, if a disease-causing mutation exists in one of the protein members in cells or tissue *in vitro* or in a patient, then a nucleic acid encoding a wild-type protein can be introduced into tissue cells of the patient. The exogenous nucleic acid can be used to replace the corresponding endogenous defective gene by, e.g., homologous recombination. *See* U.S. Patent No. 6,010,908, which is incorporated herein by reference. Alternatively, if the disease-causing mutation is a recessive mutation, the exogenous nucleic acid is simply used to express a wild-type protein in addition to the endogenous mutant protein. In another approach, the method disclosed in U.S. Patent No. 6,077,705 may be employed in gene therapy. That is, the patient is administered both a nucleic acid construct encoding a ribozyme and a nucleic acid construct comprising a ribozyme resistant gene encoding a wild type form of the gene product. As a result, undesirable expression of the endogenous gene is inhibited and a desirable wild-type exogenous gene is introduced. In yet another embodiment, if the endogenous gene is of wild-type and the level of expression of the protein encoded thereby is desired to be increased, additional copies of wild-type exogenous genes may be

introduced into the patient by gene therapy, or alternatively, a gene activation method such as that disclosed in U.S. Patent No. 5,641,670 may be used.

Various gene therapy methods are well known in the art. Successes in gene therapy have been reported recently. *See e.g.*, Kay *et al.*, *Nature Genet.*, 24:257-61 (2000); Cavazzana-Calvo *et al.*, *Science*, 288:669 (2000); and Blaese *et al.*, *Science*, 270: 475 (1995); Kantoff, *et al.*, *J. Exp. Med.* 166:219 (1987).

Any suitable gene therapy methods may be used for the purposes of the present invention. Generally, a nucleic acid encoding a desirable protein (e.g., one selected from survivin, HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT) is incorporated into a suitable expression vector and is operably linked to a promoter in the vector. Suitable promoters include but are not limited to viral transcription promoters derived from adenovirus, simian virus 40 (SV40) (e.g., the early and late promoters of SV40), Rous sarcoma virus (RSV), and cytomegalovirus (CMV) (e.g., CMV immediate-early promoter), human immunodeficiency virus (HIV) (e.g., long terminal repeat (LTR)), vaccinia virus (e.g., 7.5K promoter), and herpes simplex virus (HSV) (e.g., thymidine kinase promoter). Where tissue-specific expression of the exogenous gene is desirable, tissue-specific promoters may be operably linked to the exogenous gene. In addition, selection markers may also be included in the vector for purposes of selecting, *in vitro*, those cells that contain the exogenous gene. Various selection markers known in the art may be used including, but not limited to, e.g., genes conferring resistance to neomycin, hygromycin, zeocin, and the like.

In one embodiment, the exogenous nucleic acid (gene) is incorporated into a plasmid DNA vector. Many commercially available expression vectors may be useful for the present invention, including, e.g., pCEP4, pcDNAI, pIND, pSecTag2, pVAX1, pcDNA3.1, and pBI-EGFP, and pDisplay.

Various viral vectors may also be used. Typically, in a viral vector, the viral genome is engineered to eliminate the disease-causing capability of the virus, e.g., the ability to replicate in the host cells. The exogenous nucleic acid to be introduced into cells or tissue *in vitro* or in a patient may be incorporated into the engineered viral genome, e.g., by inserting it into a viral gene that is non-essential to the viral infectivity. Viral vectors are convenient to use as they can be easily introduced into cells, tissues and

patients by way of infection. Once in the host cell, the recombinant virus typically is integrated into the genome of the host cell. In rare instances, the recombinant virus may also replicate and remain as extrachromosomal elements.

A large number of retroviral vectors have been developed for gene therapy.

5 These include vectors derived from oncoretroviruses (e.g., MLV), lentiviruses (e.g., HIV and SIV) and other retroviruses. For example, gene therapy vectors have been developed based on murine leukemia virus (*See*, Cepko, *et al.*, *Cell*, 37:1053-1062 (1984), Cone and Mulligan, *Proc. Natl. Acad. Sci. U.S.A.*, 81:6349-6353 (1984)), mouse mammary tumor virus (*See*, Salmons *et al.*, *Biochem. Biophys. Res. Commun.*, 159:1191-1198 (1984)),
10 gibbon ape leukemia virus (*See*, Miller *et al.*, *J. Virology*, 65:2220-2224 (1991)), HIV, (*See* Shimada *et al.*, *J. Clin. Invest.*, 88:1043-1047 (1991)), and avian retroviruses (*See* Cosset *et al.*, *J. Virology*, 64:1070-1078 (1990)). In addition, various retroviral vectors are also described in U.S. Patent Nos. 6,168,916; 6,140,111; 6,096,534; 5,985,655; 5,911,983; 4,980,286; and 4,868,116, all of which are incorporated herein by reference.

15 Adeno-associated virus (AAV) vectors have been successfully tested in clinical trials. *See e.g.*, Kay *et al.*, *Nature Genet.* 24:257-61 (2000). AAV is a naturally occurring defective virus that requires other viruses such as adenoviruses or herpes viruses as helper viruses. *See* Muzyczka, *Curr. Top. Microbiol. Immun.*, 158:97 (1992). A recombinant AAV virus useful as a gene therapy vector is disclosed in U.S. Patent No.
20 6,153,436, which is incorporated herein by reference.

Adenoviral vectors can also be useful for purposes of gene therapy in accordance with the present invention. For example, U.S. Patent No. 6,001,816 discloses an adenoviral, which is used to deliver a leptin gene intravenously to a mammal to treat obesity. Other recombinant adenoviral vectors may also be used, which include those
25 disclosed in U.S. Patent Nos. 6,171,855; 6,140,087; 6,063,622; 6,033,908; and 5,932,210, and Rosenfeld *et al.*, *Science*, 252:431-434 (1991); and Rosenfeld *et al.*, *Cell*, 68:143-155 (1992).

Other useful viral vectors include recombinant hepatitis viral vectors (*See, e.g.*, U.S. Patent No. 5,981,274), and recombinant entomopox vectors (*See, e.g.*, U.S. Patent
30 Nos. 5,721,352 and 5,753,258).

Other non-traditional vectors may also be used for purposes of this invention. For example, International Publication No. WO 94/18834 discloses a method of delivering DNA into mammalian cells by conjugating the DNA to be delivered with a polyelectrolyte to form a complex. The complex may be microinjected into or taken up by cells.

The exogenous gene fragment or plasmid DNA vector containing the exogenous gene may also be introduced into cells by way of receptor-mediated endocytosis. *See e.g.*, U.S. Patent No. 6,090,619; Wu and Wu, *J. Biol. Chem.*, 263:14621 (1988); Curiel *et al.*, *Proc. Natl. Acad. Sci. USA*, 88:8850 (1991). For example, U.S. Patent No. 6,083,741 discloses introducing an exogenous nucleic acid into mammalian cells by associating the nucleic acid to a polycation moiety (e.g., poly-L-lysine having 3-100 lysine residues), which is itself coupled to an integrin receptor binding moiety (e.g., a cyclic peptide having the sequence Arg-Gly-Asp).

Alternatively, the exogenous nucleic acid or vectors containing it can also be delivered into cells via amphiphiles. *See e.g.*, U.S. Patent No. 6,071,890. Typically, the exogenous nucleic acid or a vector containing the nucleic acid forms a complex with the cationic amphiphile. Mammalian cells contacted with the complex can readily take it up.

The exogenous gene can be introduced into cells or tissue *in vitro* or in a patient for purposes of gene therapy by various methods known in the art. For example, the exogenous gene sequences alone or in a conjugated or complex form described above, or incorporated into viral or DNA vectors, may be administered directly by injection into an appropriate tissue or organ of a patient. Alternatively, catheters or like devices may be used to deliver exogenous gene sequences, complexes, or vectors into a target organ or tissue. Suitable catheters are disclosed in, e.g., U.S. Patent Nos. 4,186,745; 5,397,307; 5,547,472; 5,674,192; and 6,129,705, all of which are incorporated herein by reference.

In addition, the exogenous gene or vectors containing the gene can be introduced into isolated cells using any known techniques such as calcium phosphate precipitation, microinjection, lipofection, electroporation, biolistics, receptor-mediated endocytosis, and the like. Cells expressing the exogenous gene may be selected and redelivered back to the patient by, e.g., injection or cell transplantation. The appropriate amount of cells delivered to a patient will vary with patient conditions, and desired effect, which can be

determined by a skilled artisan. *See e.g.*, U.S. Patent Nos. 6,054,288; 6,048,524; and 6,048,729. Preferably, the cells used are autologous, i.e., cells obtained from the patient being treated.

6.3.3. Small Organic Compounds

Defective conditions or disorders in cells or tissue *in vitro* or in a patient associated with decreased concentration or activity of a survivin-containing protein complex, survivin, or a survivin-interacting protein identified in accordance with the present invention, can also be ameliorated by administering to the patient a compound identified by the methods described in Sections 5.3.1.4, 5.2, and Section 5.4, which is capable of modulating the functions of the protein complex or the survivin-interacting protein, e.g., by triggering or initiating, enhancing or stabilizing protein-protein interaction between the interacting protein members of the protein complex, or the mutant forms of such interacting protein members found in the patient.

7. Cell and Animal Models

In another aspect of the present invention, cell and animal models are provided in which one or more of the survivin-containing protein complexes identified in the present invention, or survivin itself, or a member, or members of the group consisting of HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT, exhibit aberrant function, activity, or concentration when compared with wildtype cells and animals (e.g., increased or decreased concentration, altered interactions between protein complex constituents due to mutations in interaction domains, and/or altered distribution or localization of the protein complexes or constituents thereof in organs, tissues, cells, or cellular compartments). Such cell and animal models are useful tools for studying cellular functions and biological processes associated with the protein complexes of the present invention, or with survivin itself, or with a survivin-interacting protein identified in accordance with the present invention. Such cell and animal models are also useful tools for studying disorders and diseases associated with the protein complexes of the present invention, or survivin itself, or a member, or members of the group HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT, and for testing various methods for

modulating the cellular functions, and for treating the diseases and disorders, associated with aberrations in these protein complexes or the protein constituents thereof.

7.1. Cell Models

5 Cell models having an aberrant form of one or more of the protein complexes of the present invention are provided in accordance with the present invention.

The cell models may be established by isolating, from a patient, cells having an aberrant form of one or more of the protein complexes of the present invention. The isolated cells may be cultured *in vitro* as a primary cell culture. Alternatively, the cells

10 obtained from the primary cell culture or directly from the patient may be immortalized to establish a human cell line. Any methods for constructing immortalized human cell lines may be used in this respect. *See generally* Yeager and Reddel, *Curr. Opin. Biotech.*, 10:465-469 (1999). For example, the human cells may be immortalized by transfection of plasmids expressing the SV40 early region genes (*See e.g.*, Jha *et al.*, *Exp. Cell Res.*, 245:1-7 (1998)), introduction of the HPV E6 and E7 oncogenes (*See e.g.*, Reznikoff *et al.*, *Genes Dev.*, 8:2227-2240 (1994)), and infection with Epstein-Barr virus (*See e.g.*, Tahara *et al.*, *Oncogene*, 15:1911-1920 (1997)). Alternatively, the human cells may be immortalized by recombinantly expressing the gene for the human telomerase catalytic subunit hTERT in the human cells. *See* Bodnar *et al.*, *Science*, 279:349-352

20 (1998).
In alternative embodiments, cell models are provided by recombinantly manipulating appropriate host cells. The host cells may be bacteria cells, yeast cells, insect cells, plant cells, animal cells, and the like. Preferably, the cells are derived from mammals, most preferably humans. The host cells may be obtained directly from an

25 individual, or a primary cell culture, or preferably an immortal stable human cell line. In a preferred embodiment, human embryonic stem cells or pluripotent cell lines derived from human stem cells are used as host cells. Methods for obtaining such cells are disclosed in, e.g., Shamblott, *et al.*, *Proc. Natl. Acad. Sci. USA*, 95:13726-13731 (1998) and Thomson *et al.*, *Science*, 282:1145-1147 (1998).

30 In one embodiment, a cell model is provided by recombinantly expressing one or more of the protein complexes of the present invention in cells that do not normally

express such protein complexes. For example, cells that do not contain a particular protein complex may be engineered to express the protein complex. In a specific embodiment, a particular human protein complex is expressed in non-human cells. The cell model may be prepared by introducing into host cells nucleic acids encoding all
5 interacting protein members required for the formation of a particular protein complex, and expressing the protein members in the host cells. For this purpose, the recombinant expression methods described in Section 2.2 may be used. In addition, the methods for introducing nucleic acids into host cells disclosed in the context of gene therapy in Section 6.2.2 may also be used.

10 In another embodiment, a cell model over-expressing one or more of the protein complexes of the present invention is provided. The cell model may be established by increasing the expression level of one or more of the interacting protein members of the protein complexes. In a specific embodiment, all interacting protein members of a particular protein complex are over-expressed. The over-expression may be achieved by
15 introducing into host cells exogenous nucleic acids encoding the proteins to be over-expressed, and selecting those cells that over-express the proteins. The expression of the exogenous nucleic acids may be transient or, preferably stable. The recombinant expression methods described in Section 2.2, and the methods for introducing nucleic acids into host cells disclosed in the context of gene therapy in Section 6.2.2 may be
20 used. Alternatively, the gene activation method disclosed in U.S. Patent No. 5,641,670 can be used. Any host cells may be employed for establishing the cell model. Preferably, human cells lacking a protein complex to be over-expressed, or having a normal concentration of the protein complex, are used as host cells. The host cells may be obtained directly from an individual, or a primary cell culture, or preferably a stable
25 immortal human cell line. In a preferred embodiment, human embryonic stem cells or pluripotent cell lines derived from human stem cells are used as host cells. Methods for obtaining such cells are disclosed in, e.g., Shamblott, *et al.*, *Proc. Natl. Acad. Sci. USA*, 95:13726-13731 (1998), and Thomson *et al.*, *Science*, 282:1145-1147 (1998).

30 In yet another embodiment, a cell model expressing an abnormally low level of one or more of the protein complexes of the present invention is provided. Typically, the cell model is established by genetically manipulating cells that express a normal and

detectable level of a protein complex identified in accordance with the present invention. Generally the expression level of one or more of the interacting protein members of the protein complex is reduced by recombinant methods. In a specific embodiment, the expression of all interacting protein members of a particular protein complex is reduced.

5 The reduced expression may be achieved by “knocking out” the genes encoding one or more interacting protein members. Alternatively, mutations that can cause reduced expression level (e.g., reduced transcription and/or translation efficiency, and decreased mRNA stability) may also be introduced into the gene by homologous recombination. A gene encoding a ribozyme or antisense compound specific to the mRNA encoding an
10 interacting protein member may also be introduced into the host cells, preferably stably integrated into the genome of the host cells. In addition, a gene encoding an antibody or fragment thereof specific to an interacting protein member may also be introduced into the host cells. The recombinant expression methods described in Sections 2.2, 6.1 and 6.2 can all be used for purposes of manipulating the host cells.

15 The present invention also contemplates a cell model provided by recombinant DNA techniques that exhibits aberrant interactions between the interacting protein members of a protein complex identified in the present invention. For example, variants of the interacting protein members of a particular protein complex exhibiting altered protein-protein interaction properties and the nucleic acid variants encoding such variant
20 proteins may be obtained by random or site-directed mutagenesis in combination with a protein-protein interaction assay system, particularly the yeast two-hybrid system described in Section 5.3.1. Essentially, the genes encoding one or more interacting protein members of a particular protein complex may be subject to random or site-specific mutagenesis and the mutated gene sequences are used in yeast two-hybrid system
25 to test the protein-protein interaction characteristics of the protein variants encoded by the gene variants. In this manner, variants of the interacting protein members of the protein complex may be identified that exhibit altered protein-protein interaction properties in forming the protein complex, e.g., increased or decreased binding affinity, and the like. The nucleic acid variants encoding such protein variants may be introduced into host
30 cells by the methods described above, preferably into host cells that normally do not express the interacting proteins.

7.2. Cell-Based Assays

The cell models of the present invention containing an aberrant form of a survivin-containing protein complex of the present invention are useful in screening assays for identifying compounds useful in treating diseases and disorders involving apoptosis such as cancer, viral infection, AIDS, ischemia, stroke, autoimmune diseases, neurodegenerative diseases, inflammatory disorders, sepsis, and osteoporosis. In addition, they may also be used in *in vitro* pre-clinical assays for testing compounds, such as those identified in the screening assays of the present invention.

For example, cells may be treated with compounds to be tested and assayed for the compound's activity. A variety of parameters relevant to particularly physiological disorders or diseases may be analyzed.

7.3. Transgenic Animals

In another aspect of the present invention, transgenic non-human animals are created expressing an aberrant form of one or more of the survivin-containing protein complexes of the present invention. Animals of any species may be used to generate the transgenic animal models, including but not limited to, mice, rats, hamsters, sheep, pigs, rabbits, guinea pigs, preferably non-human primates such as monkeys, chimpanzees, baboons, and the like.

In one embodiment, transgenic animals are made to over-express one or more protein complexes formed from survivin, or a derivative, fragment or homologue thereof (including the animal counterpart of survivin, i.e., an orthologue) and a member, or members, of the group of survivin-interacting proteins including HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT, or derivatives, fragments or homologues thereof (including orthologues). Over-expression may be directed in a tissue or cell type that normally expresses animal counterparts of such protein complexes. Consequently, the concentration of the protein complex(es) will be elevated to higher levels than normal. Alternatively, the one or more protein complexes are expressed in tissues or cells that do not normally express such proteins and hence do not normally contain the protein complexes of the present invention. In a specific embodiment, human

survivin and a human protein, or proteins, from the group of survivin-interacting proteins including HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT, are expressed in the transgenic animals.

To achieve over-expression in transgenic animals, the transgenic animals are made such that they contain and express exogenous, orthologous genes encoding survivin or a homologue, derivative or mutant form thereof and one or more survivin-interacting proteins or homologues, derivatives or mutant forms thereof. Preferably, the exogenous genes are human genes. Such exogenous genes may be operably linked to a native or non-native promoter, preferably a non-native promoter. For example, an exogenous survivin gene may be operably linked to a promoter that is not the native survivin promoter. If the expression of the exogenous gene is desired to be limited to a particular tissue, an appropriate tissue-specific promoter may be used.

Over-expression may also be achieved by manipulating the native promoter to create mutations that lead to gene over-expression, or by a gene activation method such as that disclosed in U.S. Patent No. 5,641,670 as described above.

In another embodiment, the transgenic animal expresses an abnormally low concentration of the complex comprising survivin and one or more of the survivin-interacting proteins from the group HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT. In a specific embodiment, the transgenic animal is a "knockout" animal wherein the endogenous gene encoding the animal orthologue of survivin and/or an endogenous gene encoding an animal orthologue of a survivin-interacting protein are knocked out. In a specific embodiment, the expression of the animal orthologues of both survivin and a survivin-interacting protein, or proteins, from the group HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT are reduced or knocked out.

The reduced expression may be achieved by knocking out the genes encoding one or both interacting protein members, typically by homologous recombination. Alternatively, mutations that can cause reduced expression (e.g., reduced transcription and/or translation efficiency, or decreased mRNA stability) may also be introduced into the endogenous genes by homologous recombination. Genes encoding ribozymes or antisense compounds specific to the mRNAs encoding the interacting protein members may also be introduced into the transgenic animal. In addition, genes encoding

antibodies or fragments thereof specific to the interacting protein members may also be introduced into the transgenic animal.

In an alternate embodiment, transgenic animals are made in which the endogenous genes encoding the animal orthologues of survivin and one or more survivin-
5 interacting proteins from the group HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT are replaced with orthologous human genes.

In yet another embodiment, the transgenic animal of this invention expresses specific mutant forms of survivin and one or more survivin-interacting proteins from the group HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT that
10 exhibit aberrant interactions. For this purpose, variants of survivin and one or more survivin-interacting proteins from the group HDLC1, beta-actin, DNA helicase II, COPP, OSTP, SLC8A1, A2-CAT exhibiting altered protein-protein interaction properties, and the nucleic acid variants encoding such variant proteins, may be obtained by random or site-specific mutagenesis in combination with a protein-protein interaction assay system,
15 particularly the yeast two-hybrid system described in Section 5.3.1. For example, variants of survivin and HDLC1 exhibiting increased, decreased or abolished binding affinity to each other may be identified and isolated. The transgenic animal of the present invention may be made to express such protein variants by modifying the endogenous genes. Alternatively, the nucleic acid variants may be introduced exogenously into the
20 transgenic animal genome to express the protein variants therein. In a specific embodiment, the exogenous nucleic acid variants are derived from orthologous human genes and the corresponding endogenous genes are knocked out.

Any techniques known in the art for making transgenic animals may be used for purposes of the present invention. For example, the transgenic animals of the present
25 invention may be provided by methods described in, e.g., Jaenisch, *Science*, 240:1468-1474 (1988); Capecchi, *et al.*, *Science*, 244:1288-1291 (1989); Hasty *et al.*, *Nature*, 350:243 (1991); Shinkai *et al.*, *Cell*, 68:855 (1992); Mombaerts *et al.*, *Cell*, 68:869 (1992); Philpott *et al.*, *Science*, 256:1448 (1992); Snouwaert *et al.*, *Science*, 257:1083 (1992); Donehower *et al.*, *Nature*, 356:215 (1992); Hogan *et al.*, *Manipulating the Mouse Embryo; A Laboratory Manual*, 2nd edition, Cold Spring Harbor Laboratory Press, 1994;
30

and U.S. Patent Nos. 4,873,191; 5,800,998; 5,891,628, all of which are incorporated herein by reference.

Generally, the founder lines may be established by introducing appropriate exogenous nucleic acids into, or modifying an endogenous gene in, germ lines, embryonic stem cells, embryos, or sperm which are then used in producing a transgenic animal. The gene introduction may be conducted by various methods including those described in Sections 2.2, 6.1 and 6.2. *See also*, Van der Putten *et al.*, *Proc. Natl. Acad. Sci. USA*, 82:6148-6152 (1985); Thompson *et al.*, *Cell*, 56:313-321 (1989); Lo, *Mol. Cell. Biol.*, 3:1803-1814 (1983); Gordon, *Transgenic Animals*, *Intl. Rev. Cytol.* 115:171-229 (1989); and Lavitrano *et al.*, *Cell*, 57:717-723 (1989). In a specific embodiment, the exogenous gene is incorporated into an appropriate vector, such as those described in Sections 2.2 and 6.2, and is transformed into embryonic stem (ES) cells. The transformed ES cells are then injected into a blastocyst. The blastocyst with the transformed ES cells is then implanted into a surrogate mother animal. In this manner, a chimeric founder line animal containing the exogenous nucleic acid (transgene) may be produced.

Preferably, site-specific recombination is employed to integrate the exogenous gene into a specific predetermined site in the animal genome, or to replace an endogenous gene or a portion thereof with the exogenous sequence. Various site-specific recombination systems may be used including those disclosed in Sauer, *Curr. Opin. Biotechnol.*, 5:521-527 (1994); Capecchi, *et al.*, *Science*, 244:1288-1291 (1989); and Gu *et al.*, *Science*, 265:103-106 (1994). Specifically, the Cre/lox site-specific recombination system known in the art may be conveniently used which employs the bacteriophage P1 protein Cre recombinase and its recognition sequence *loxP*. *See* Rajewsky *et al.*, *J. Clin. Invest.*, 98:600-603 (1996); Sauer, *Methods*, 14:381-392 (1998); Gu *et al.*, *Cell*, 73:1155-1164 (1993); Araki *et al.*, *Proc. Natl. Acad. Sci. USA*, 92:160-164 (1995); Lakso *et al.*, *Proc. Natl. Acad. Sci. USA*, 89:6232-6236 (1992); and Orban *et al.*, *Proc. Natl. Acad. Sci. USA*, 89:6861-6865 (1992).

The transgenic animals of the present invention may be transgenic animals that carry a transgene in all cells or mosaic transgenic animals carrying a transgene only in

certain cells, e.g., somatic cells. The transgenic animals may have a single copy or multiple copies of a particular transgene.

The founder transgenic animals thus produced may be bred to produce various offsprings. For example, they can be inbred, outbred, and crossbred to establish
5 homozygous lines, heterozygous lines, and compound homozygous or heterozygous lines.

8. Pharmaceutical Compositions and Formulations

For example, such therapeutic agents include, but are not limited to, (1) small
10 organic compounds selected based on the screening methods of the present invention capable of interfering with the interaction between survivin and an interactor thereof, (2) antisense compounds specifically hybridizable to survivin nucleic acids (gene or mRNA) (3) antisense compounds specific to the gene or mRNA of a survivin interactor, (4) ribozyme compounds specific to survivin nucleic acids (gene or mRNA), (5) ribozyme
15 compounds specific to the gene or mRNA of a survivin interactor, (6) antibodies immunoreactive with survivin or a survivin interactor, (7) antibodies selectively immunoreactive with a protein complex of the present invention, (8) small organic compounds capable of binding a protein complex of the present invention, (9) small peptide compounds as described above (optionally linked to a transporter) capable of
20 interacting with survivin or a survivin interactor, (10) nucleic acids encoding the antibodies or peptides, etc.

The compositions are prepared as a pharmaceutical formulation suitable for administration into a patient. Accordingly, the present invention also extends to pharmaceutical compositions, medicaments, drugs or other compositions containing one
25 or more of the therapeutic agent in accordance with the present invention.

In the pharmaceutical composition, an active compound identified in accordance with the present invention can be in any pharmaceutically acceptable salt form. As used herein, the term "pharmaceutically acceptable salts" refers to the relatively non-toxic, organic or inorganic salts of the compounds of the present invention, including inorganic
30 or organic acid addition salts of the compound. Examples of such salts include, but are not limited to, hydrochloride salts, sulfate salts, bisulfate salts, borate salts, nitrate salts,

acetate salts, phosphate salts, hydrobromide salts, laurylsulfonate salts, glucoheptonate salts, oxalate salts, oleate salts, laurate salts, stearate salts, palmitate salts, valerate salts, benzoate salts, naththylate salts, mesylate salts, tosylate salts, citrate salts, lactate salts, maleate salts, succinate salts, tartrate salts, fumarate salts, and the like. See, e.g., Berge, et al., J. Pharm. Sci., 66:1-19 (1977).

For oral delivery, the active compounds can be incorporated into a formulation that includes pharmaceutically acceptable carriers such as binders (e.g., gelatin, cellulose, gum tragacanth), excipients (e.g., starch, lactose), lubricants (e.g., magnesium stearate, silicon dioxide), disintegrating agents (e.g., alginate, Primogel, and corn starch), and sweetening or flavoring agents (e.g., glucose, sucrose, saccharin, methyl salicylate, and peppermint). The formulation can be orally delivered in the form of enclosed gelatin capsules or compressed tablets. Capsules and tablets can be prepared in any conventional techniques. The capsules and tablets can also be coated with various coatings known in the art to modify the flavors, tastes, colors, and shapes of the capsules and tablets. In addition, liquid carriers such as fatty oil can also be included in capsules.

Suitable oral formulations can also be in the form of suspension, syrup, chewing gum, wafer, elixir, and the like. If desired, conventional agents for modifying flavors, tastes, colors, and shapes of the special forms can also be included. In addition, for convenient administration by enteral feeding tube in patients unable to swallow, the active compounds can be dissolved in an acceptable lipophilic vegetable oil vehicle such as olive oil, corn oil and safflower oil.

The active compounds can also be administered parenterally in the form of solution or suspension, or in lyophilized form capable of conversion into a solution or suspension form before use. In such formulations, diluents or pharmaceutically acceptable carriers such as sterile water and physiological saline buffer can be used. Other conventional solvents, pH buffers, stabilizers, anti-bacterial agents, surfactants, and antioxidants can all be included. For example, useful components include sodium chloride, acetate, citrate or phosphate buffers, glycerin, dextrose, fixed oils, methyl parabens, polyethylene glycol, propylene glycol, sodium bisulfate, benzyl alcohol, ascorbic acid, and the like. The parenteral formulations can be stored in any conventional containers such as vials and ampoules.

Routes of topical administration include nasal, bucal, mucosal, rectal, or vaginal applications. For topical administration, the active compounds can be formulated into lotions, creams, ointments, gels, powders, pastes, sprays, suspensions, drops and aerosols. Thus, one or more thickening agents, humectants, and stabilizing agents can be included in the formulations. Examples of such agents include, but are not limited to, polyethylene glycol, sorbitol, xanthan gum, petrolatum, beeswax, or mineral oil, lanolin, squalene, and the like. A special form of topical administration is delivery by a transdermal patch. Methods for preparing transdermal patches are disclosed, e.g., in Brown, *et al.*, *Annual Review of Medicine*, 39:221-229 (1988), which is incorporated herein by reference.

Subcutaneous implantation for sustained release of the active compounds may also be a suitable route of administration. This entails surgical procedures for implanting an active compound in any suitable formulation into a subcutaneous space, e.g., beneath the anterior abdominal wall. *See, e.g.*, Wilson *et al.*, *J. Clin. Psych.* 45:242-247 (1984). Hydrogels can be used as a carrier for the sustained release of the active compounds. Hydrogels are generally known in the art. They are typically made by crosslinking high molecular weight biocompatible polymers into a network that swells in water to form a gel like material. Preferably, hydrogels is biodegradable or biosorbable. For purposes of this invention, hydrogels made of polyethylene glycols, collagen, or poly(glycolic-co-L-lactic acid) may be useful. *See, e.g.*, Phillips *et al.*, *J. Pharmaceut. Sci.* 73:1718-1720 (1984).

The active compounds can also be conjugated, to a water soluble non-immunogenic non-peptidic high molecular weight polymer to form a polymer conjugate. For example, an active compound is covalently linked to polyethylene glycol to form a conjugate. Typically, such a conjugate exhibits improved solubility, stability, and reduced toxicity and immunogenicity. Thus, when administered to a patient, the active compound in the conjugate can have a longer half-life in the body, and exhibit better efficacy. *See generally*, Burnham, *Am. J. Hosp. Pharm.*, 15:210-218 (1994). PEGylated proteins are currently being used in protein replacement therapies and for other therapeutic uses. For example, PEGylated interferon (PEG-INTRON A[®]) is clinically used for treating Hepatitis B. PEGylated adenosine deaminase (ADAGEN[®]) is being

used to treat severe combined immunodeficiency disease (SCIDS). PEGylated L-asparaginase (ONCAPSPAR[®]) is being used to treat acute lymphoblastic leukemia (ALL). It is preferred that the covalent linkage between the polymer and the active compound and/or the polymer itself is hydrolytically degradable under physiological conditions. Such conjugates known as “prodrugs” can readily release the active compound inside the body. Controlled release of an active compound can also be achieved by incorporating the active ingredient into microcapsules, nanocapsules, or hydrogels generally known in the art.

Liposomes can also be used as carriers for the active compounds of the present invention. Liposomes are micelles made of various lipids such as cholesterol, phospholipids, fatty acids, and derivatives thereof. Various modified lipids can also be used. Liposomes can reduce the toxicity of the active compounds, and increase their stability. Methods for preparing liposomal suspensions containing active ingredients therein are generally known in the art. *See, e.g.*, U.S. Patent No. 4,522,811; Prescott, Ed., *Methods in Cell Biology*, Volume XIV, Academic Press, New York, N.Y. (1976).

The active compounds can also be administered in combination with another active agent that synergistically treats or prevents the same symptoms or is effective for another disease or symptom in the patient treated so long as the other active agent does not interfere with or adversely affect the effects of the active compounds of this invention. Such other active agents include but are not limited to anti-inflammation agents, antiviral agents, antibiotics, antifungal agents, antithrombotic agents, cardiovascular drugs, cholesterol lowering agents, anti-cancer drugs, hypertension drugs, and the like.

Generally, the toxicity profile and therapeutic efficacy of the therapeutic agents can be determined by standard pharmaceutical procedures in cell models or animal models, e.g., those provided in Section 7. As is known in the art, the LD₅₀ represents the dose lethal to about 50% of a tested population. The ED₅₀ is a parameter indicating the dose therapeutically effective in about 50% of a tested population. Both LD₅₀ and ED₅₀ can be determined in cell models and animal models. In addition, the IC₅₀ may also be obtained in cell models and animal models, which stands for the circulating plasma concentration that is effective in achieving about 50% of the maximal inhibition of the

symptoms of a disease or disorder. Such data may be used in designing a dosage range for clinical trials in humans. Typically, as will be apparent to skilled artisans, the dosage range for human use should be designed such that the range centers around the ED₅₀ and/or IC₅₀, but significantly below the LD₅₀ obtained from cell or animal models.

It will be apparent to skilled artisans that therapeutically effective amount for each active compound to be included in a pharmaceutical composition of the present invention can vary with factors including but not limited to the activity of the compound used, stability of the active compound in the patient's body, the severity of the conditions to be alleviated, the total weight of the patient treated, the route of administration, the ease of absorption, distribution, and excretion of the active compound by the body, the age and sensitivity of the patient to be treated, and the like. The amount of administration can also be adjusted as the various factors change over time.

EXAMPLES

1. Yeast Two-Hybrid System

The principles and methods of the yeast two-hybrid system have been described in detail in *The Yeast Two-Hybrid System*, Bartel and Fields, eds., pages 183-196, Oxford University Press, New York, NY, 1997. The following is thus a description of the particular procedure that we used, which was applied to all proteins.

The cDNA encoding the bait protein was generated by PCR from cDNA prepared from a desired tissue. The cDNA product was then introduced by recombination into the yeast expression vector pGBT.Q, which is a close derivative of pGBT.C (*See Bartel et al., Nat Genet.*, 12:72-77 (1996)) in which the polylinker site has been modified to include M13 sequencing sites. The new construct was selected directly in the yeast strain PNY200 for its ability to drive tryptophane synthesis (genotype of this strain: *MAT α trp1-901 leu2-3,112 ura3-52 his3-200 ade2 gal4 Δ gal80*). In these yeast cells, the bait was produced as a C-terminal fusion protein with the DNA binding domain of the transcription factor Gal4 (amino acids 1 to 147). Prey libraries were transformed into the yeast strain BK100 (genotype of this strain: *MAT α trp1-901 leu2-3,112 ura3-52 his3-200 gal4 Δ gal80 LYS2::GAL-HIS3 GAL2-ADE2 met2::GAL7-lacZ*), and selected for the ability to drive leucine synthesis. In these yeast cells, each cDNA was expressed as a

fusion protein with the transcription activation domain of the transcription factor Gal4 (amino acids 768 to 881) and a 9 amino acid hemagglutinin epitope tag. PNY200 cells (MAT α mating type), expressing the bait, were then mated with BK100 cells (MATa mating type), expressing prey proteins from a prey library. The resulting diploid yeast cells expressing proteins interacting with the bait protein were selected for the ability to synthesize tryptophan, leucine, histidine, and adenine. DNA was prepared from each clone, transformed by electroporation into *E. coli* strain KC8 (Clontech KC8 electrocompetent cells, Catalog No. C2023-1), and the cells were selected on ampicillin-containing plates in the absence of either tryptophane (selection for the bait plasmid) or leucine (selection for the library plasmid). DNA for both plasmids was prepared and sequenced by the dideoxynucleotide chain termination method. The identity of the bait cDNA insert was confirmed and the cDNA insert from the prey library plasmid was identified using the BLAST program to search against public nucleotide and protein databases. Plasmids from the prey library were then individually transformed into yeast cells together with a plasmid driving the synthesis of lamin and 5 other test proteins, respectively, fused to the Gal4 DNA binding domain. Clones that gave a positive signal in the β -galactosidase assay were considered false-positives and discarded. Plasmids for the remaining clones were transformed into yeast cells together with the original bait plasmid. Clones that gave a positive signal in the β -galactosidase assay were considered true positives.

Bait sequences indicated in Table I were used in the yeast two-hybrid system described above. The isolated prey sequences are summarized in Table I. The GenBank Accession Nos. for the bait and prey proteins are also provided in Table I, upon which the bait and prey sequences are aligned.

2. Production of Antibodies Selectively Immunoreactive with Protein Complex

The survivin-interacting region of HDLC1 and the HDLC1-interacting region of survivin are indicated in Table I in Section 2. Both regions, or fragments thereof, are recombinantly-expressed in *E. coli*. and isolated and purified. Mixing the two purified interacting regions forms a protein complex. A protein complex is also formed by mixing recombinantly expressed intact complete survivin and HDLC1. The two protein

complexes are used as antigens in immunizing a mouse. mRNA is isolated from the immunized mouse spleen cells, and first-strand cDNA is synthesized using the mRNA as a template. The V_H and V_K genes are amplified from the thus synthesized cDNAs by PCR using appropriate primers.

5 The amplified V_H and V_K genes are ligated together and subcloned into a phagemid vector for the construction of a phage display library. *E. coli* cells are transformed with the ligation mixtures, and thus a phage display library is established. Alternatively, the ligated V_H and V_K genes are subcloned into a vector suitable for ribosome display in which the V_H-V_K sequence is under the control of a T7 promoter. See
10 Schaffitzel *et al.*, *J. Immun. Meth.*, 231:119-135 (1999).

 The libraries are screened for their ability to bind survivin-HDLC1 complex and survivin or HDLC1, alone. Several rounds of screening are generally performed. Clones corresponding to scFv fragments that bind the survivin-HDLC1 complex, but not isolated survivin or HDLC1 are selected and purified. A single purified clone is used to prepare
15 an antibody selectively immunoreactive with the complex comprising survivin and HDLC1. The antibody is then verified by an immunochemistry method such as RIA and ELISA.

 In addition, the clones corresponding to scFv fragments that bind the complex comprising survivin and HDLC1, and also bind isolated survivin and/or HDLC1 may be
20 selected. The scFv genes in the clones are diversified by mutagenesis methods such as oligonucleotide-directed mutagenesis, error-prone PCR (See Lin-Goerke *et al.*, *Biotechniques*, 23:409 (1997)), dNTP analogues (See Zaccole *et al.*, *J. Mol. Biol.*, 255:589 (1996)), and other methods. The diversified clones are further screened in phage display or ribosome display libraries. In this manner, scFv fragments selectively
25 immunoreactive with the complex comprising survivin and HDLC1 may be obtained.

3. Yeast Screen To Identify Small Molecule Inhibitors Of The Interaction Between survivin and HDLC1

 Beta-galactosidase is used as a reporter enzyme to signal the interaction between
30 yeast two-hybrid protein pairs expressed from plasmids in *Saccharomyces cerevisiae*. Yeast strain MY209 (*ade2 his3 leu2 trp1 cyh2 ura3::GAL1p-lacZ gal4 gal80*

lys2::GAL1p-HIS3) bearing one plasmid with the genotype of *LEU2 CEN4 ARS1*
ADH1p-SV40NLS-GAL4 (768-881)-HDLC1-PGK1t AmpR ColE1_ori, and another
plasmid having a genotype of *TRP1 CEN4 ARS ADH1p-GAL4(1-147)-survivin-ADH1t*
AmpR ColE1_ori is cultured in synthetic complete media lacking leucine and tryptophan
5 (SC –Leu –Trp) overnight at 30°C. The survivin and HDLC1 nucleic acids in the
plasmids can code for the full-length survivin and HDLC1 proteins, respectively, or
fragments thereof. This culture is diluted to 0.01 OD₆₃₀ units/ml using SC –Leu –Trp
media. The diluted MY209 culture is dispensed into 96-well microplates. Compounds
from a library of small molecules are added to the microplates; the final concentration of
10 test compounds is approximately 60µM. The assay plates are incubated at 30°C
overnight.

The following day an aliquot of concentrated substrate/lysis buffer is added to
each well and the plates incubated at 37°C for 1-2 hours. At an appropriate time an
aliquot of stop solution is added to each well to halt the beta-galactosidase reaction. For
15 all microplates an absorbance reading is obtained to assay the generation of product from
the enzyme substrate. The presence of putative inhibitors of the interaction between
survivin and HDLC1 results in inhibition of the beta-galactosidase signal generated by
MY209. Additional testing eliminates compounds that decreased expression of beta-
galactosidase by affecting yeast cell growth and non-specific inhibitors that affected the
20 beta-galactosidase signal generated by the interaction of an unrelated protein pair.

Once a hit, i.e., a compound which inhibits the interaction between the viral and
cellular proteins, is obtained, the compound is identified and subjected to further testing
wherein the compounds are assayed at several concentrations to determine an IC₅₀ value,
this being the concentration of the compound at which the signal seen in the two-hybrid
25 assay described in this Example is 50% of the signal seen in the absence of the inhibitor.

All publications and patent applications mentioned in the specification are
indicative of the level of those skilled in the art to which this invention pertains. All
publications and patent applications are herein incorporated by reference to the same
extent as if each individual publication or patent application was specifically and
30 individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.